

Design and Development of New Zinc Porphyrin  
Molecules as High Performing Photoacoustic  
Contrast Agents

AMINA YASIN

Doctor of Philosophy

UNIVERSITI MALAYSIA PAHANG



## **SUPERVISOR'S DECLARATION**

We hereby declare that we have checked this thesis and in our opinion, this thesis is adequate in terms of scope and quality for the award of the degree of Doctor of Philosophy in Chemistry.

---

(Supervisor's Signature)

Full Name : PROFESSOR DATO' DR. MASHITAH MOHD YUSOFF  
Position : PROFESSOR, DEPUTY VICE CHANCELLOR UMP  
Date : 10-06-2018

---

(Co-supervisor's Signature)

Full Name : PROFESSOR DR. JOSE RAJAN  
Position : PROFESSOR, DEAN OF RESEARCH UMP  
Date : 10-06-2018

---

(Co-supervisor's Signature)

Full Name : DR. MD SHAHEEN SARKAR  
Position : SENIOR LECTURER  
Date : 10-06-2018



## **STUDENT'S DECLARATION**

I hereby declare that the work in this thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

---

(Student's Signature)

Full Name : AMINA YASIN

ID Number : PSK13002

Date : 10-06-2018

Design and Development of New Zinc Porphyrin Molecules as High Performing  
Photoacoustic Contrast Agents

AMINA YASIN

Thesis submitted in fulfillment of the requirements  
for the award of the degree of  
Doctor of Philosophy

Faculty of Industrial Sciences & Technology

UNIVERSITI MALAYSIA PAHANG

JUNE 2018

## DEDICATION

*“This thesis is dedicated to all my Teachers and Supervisors, especially Dr. Munawar Ali Munawar & Prof. Dr. Jose Rajan for providing me a source of inspiration and their countless effort, to my Father Muhammad Yasin Bhatti for his love and giving me encourage in every step of life and to my Husband Muhammad Wasif Nabeel for his patience and moral support”*

## ACKNOWLEDGEMENTS

I am obliged and would like to express my sincere gratitude to my supervisors Professor Dato' Dr. Mashitah Mohd Yusoff; Professor Dr. Jose Rajan; and Dr. Md Shaheen Sarkar for their indispensable guidance, interminable encouragement and relentless support in making this research possible. I really appreciate their supervision from the beginning to the final level that empowered me to develop an understanding of this research thoroughly. Without their advice and kind assistance, it would be a lot tougher to accomplish. I also sincerely thanks for the time spent proofreading and correcting my mistakes.

I also would like to express very special thanks to Gaussian technical support team for their suggestions and co-operation. A special appreciation should be given to Fernando R. Clemente from Gaussian technical support team whom which gave me a brand new perception about my research.

Besides my advisors, I would like to thank Dr. Chandra S. Yelleswarapu from Department of Physics, University of Massachusetts Boston, United States and Dr. Iman S. Roqan from Physical Science and Engineering Division, King Abdullah University of Science and Technology, Kingdom of Saudia Arabia for their collaboration on this project that make us possible to complete this doctoral dissertation.

My sincere thanks go to all lecturers and members of the staff of the FSTI Department, UMP, who helped me in many ways and made my education journey at UMP pleasant and unforgettable. Many thanks go to Central Laboratory, Institute of Postgraduate Studies & International Office for their co-operation, and supports during this study. This experience will be remembered as important memory for me to face the new chapter of life.

I acknowledge my sincere indebtedness and gratitude to my parents for their love, dream and sacrifice throughout my life. I am thankful for their sacrifice, patience, and understanding that were inevitable to make this work possible. Their sacrifice had inspired me from the day I learned how to read and write until what I have become now. I cannot find the appropriate words that could properly describe my appreciation for their devotion, support and faith in my ability to achieve my dreams.

I am very grateful to my dearest husband who supported me in every possible way to fulfil my dream. Without his patience, I would not be able to complete my doctoral research. My deepest gratitude to my siblings and my family for the continuous prayers, encouragement and support, at every stage of my life. I also want to share this accomplishment with my friends for always being there for me.

Lastly, I would like to thanks any person, which contributes to my dissertation project directly or indirectly. I would like to acknowledge their comments and suggestions, which was crucial for the successful completion of this research.

AMINA YASIN, June 2018

## ABSTRAK

Penyelidikan doktoral ini bertujuan untuk merekabentuk dan membangunkan molekul porfirin baru sebagai agen kontras dalam pengimejan fotoakustik (PA) seperti tomografi PA dan mikroskopi PA. Untuk kejayaan pelaksanaan teknik pengimejan PA, agen kontras yang mempamerkan koefisien PA yang besar adalah salah satu keperluan penting. Banyak bahan, sama ada biomolekul endogen seperti hemoglobin, deoxyhaemoglobin, melanin atau bahan-bahan eksogen seperti molekul organik, kluster atau nanokristal didapati menunjukkan kesan PA. Walau bagaimanapun, ia sama ada dihad oleh koefisien PA yang lemah, kedalaman penembusan yang lemah atau ketoksikan. Pengembangan molekul PA baru adalah terhad kerana faktor-faktor yang mempengaruhi kesan PA jarang diteliti secara sistematik, mengakibatkan kekurangan panduan untuk reka bentuk molekul agen kontras PA. Dalam tesis ini, hipotesis bahawa kesan PA yang besar boleh dicapai jika getaran kumpulan molekul atau berfungsi terletak di atas atom yang membentuk orbital molekul tidak dihuni paling rendah dapat dimanipulasi sedemikian rupa sehingga kekuatan pengayun dari getaran tersebut meningkat pada pengubahsuaian kimia yang sesuai, maka sebahagian besar foton yang teruja dapat disejukkan melalui pelepasan fonon. Dengan menggunakan Zn(II)porfirin sebagai sistem model, sebilangan besar molekul dimodelkan menggunakan teori berfungsi ketumpatan (DFT) semasa berdasarkan andaian di atas dan sifat asas dan keadaan teruja dikaji secara sistematik dengan menggunakan kaedah hibrid tiga parameter Becke dengan gradien korelasi berfungsi (B3LYP) Lee, Yang, dan Parr dan teras-teras berpotensi diperbetulkan yang berkesan menggunakan LANL2DZ dan 6-31G(d) dalam pakej program Gaussian 09W. Telah diperhatikan bahawa kumpulan-kumpulan berfungsi dalam kumpulan fenil kurang resonans yang melekat pada teras porfirin menguatkan kekuatan ayunan kelompok alkoksi – variasi sistematik dalam koefisien penyerapan mod C-H vibrational juga diperhatikan dengan perubahan relatif dalam resonans kumpulan molekul berlabuh. Dipengaruhi pemerhatian tersebut, empat molekul Zn(II)porfirin ( $RJ-C_n-MY_m$ , di mana  $n = 12$  atau  $8$  dan  $m = 1-4$ ) telah disintesis dengan hasil yang baik melalui pengoksidaan berasaskan asid, pemeluwapan Knoevenagel, pemeluwapan MacDonald (2+2), brominasi, logam, nyahlindung dan reaksi gandingan Sonogashira. Struktur molekul dikaji menggunakan spektroskopi NMR, MS, dan FTIR manakala ciri-ciri optik dikaji dengan kaedah spektroskopi penyerapan UV-Vis, spektroskopi fotopendarkilau dan pengiraan foton tunggal yang berkorelasi; sementara sifat-sifat PA dikaji dengan kaedah spektroskopi fotoakustik. Koefisien penyerapan menurun mengikut susunan  $RJ-C_{12}-MY_1 > RJ-C_{12}-MY_2 > RJ-C_{12}-MY_3 > RJ-C_8-MY_4$ ; oleh itu variasi yang serupa dalam isyarat PA telah diramalkan. Ramalan teori telah disahkan oleh ukuran koefisien PA dengan tomografi PA. Koefisien PA dan keamatan tomografi menurun mengikut susunan  $RJ-C_{12}-MY_1 > RJ-C_{12}-MY_2 > RJ-C_{12}-MY_3 > RJ-C_8-MY_4$ , seperti yang diramalkan oleh DFT. Pengelasan molekul terperinci dalam menentukan ciri-ciri optoelektronik seperti kepupusan koefisien molar keadaan asas, kepupusan koefisien molar keadaan teruja, hasil kuantum pendarfluor dan masa hayat keadaan teruja molekul Zn(II)porfirin yang baru disintesis diperiksa bagi menunjukkan korelasi positif antara tahap tenaga getaran, koefisien kepupusan molar, keadaan pemancaran fonon dan kesan PA. Koefisien PA yang besar diperhatikan bagi  $RJ-C_{12}-MY_1$  memberikan peningkatan PA sebanyak  $\sim 7$  kali ganda berbanding dengan ZnTPP yang digunakan sebagai rujukan adalah lebih tinggi daripada molekul kecil sedia ada. Di samping menawarkan molekul unggul untuk tomografi PA tesis dan kriteria yang diterima pakai akan membolehkan rekabentuk molekul yang lebih mudah dengan ciri-ciri fotoakustik superior dan lain-lain ciri optik tidak linear.

## ABSTRACT

This doctoral research aims to design and develop new porphyrin molecules as contrast agents in photoacoustic (PA) imaging (PAI) such as PA tomography and PA microscopy. For successful implementation of PAI techniques, contrast agents exhibiting large PA coefficients are one of the essential requirements. Many materials, either endogenous biomolecules such as haemoglobin, deoxyhaemoglobin, melanin or exogenous materials such as organic molecules, clusters or nanocrystals are shown to exhibit PA effects. However, they are either limited by poor PA coefficients, poor penetration depths or toxicity. Development of new PA molecules is limited because the factors influencing the PA effect have rarely been systematically studied, resulting in a lack of guidance for the design of molecular PA contrast agents. In this thesis, it is hypothesized that large PA effects could be achieved if vibrations of the molecular or functional groups lying above the atoms composing the lowest unoccupied molecular orbitals could be manipulated in such a way that the oscillator strength of those vibrations increase upon suitable chemical modification, then a large portion of the excited photons can be thermalized via phonon emission. By using Zn(II)porphyrin as a model system, a large number of molecules were modelled using state of the art density functional theory (DFT) based on the above assumption and their ground and excited state properties were studied systematically by using the Becke's three-parameter hybrid method with the Lee, Yang, and Parr gradient corrected correlation functional (B3LYP) with the effective core potential employing basis sets of LANL2DZ and 6-31G(d) using the Gaussian 09W program package. It was observed that less resonating functional groups in a phenyl group attached to the porphyrin core amplify the oscillation strengths of the alkoxy groups – a systematic variation in the absorption coefficients of the C-H vibrational modes was observed with relative change in the resonance of the molecular groups anchored. Motivated from this result, representative four Zn(II)porphyrin molecules (RJ-C<sub>n</sub>-MY<sub>m</sub>, where n = 12 or 8 and m = 1 – 4) were synthesized in good yields by acid-catalyzed oxidation, Knoevenagel condensation, MacDonald (2+2) condensation, bromination, metalation, deprotection, and Sonogashira coupling reactions. Structure of the molecules were studied by NMR, Mass Spectrometry, and FTIR spectroscopy; optical properties were studied by UV-Vis absorption spectroscopy, photoluminescence spectroscopy, and time-correlated single photon counting; and PA properties were studied by photoacoustic spectroscopy. The absorption coefficient decreased in the order of RJ-C<sub>12</sub>-MY<sub>1</sub> > RJ-C<sub>12</sub>-MY<sub>2</sub> > RJ-C<sub>12</sub>-MY<sub>3</sub> > RJ-C<sub>8</sub>-MY<sub>4</sub>; and therefore, a similar variation in the PA signals was predicted. Theoretical predictions were validated by PA coefficients measurements and PA tomography. The PA coefficients and tomographic intensities decreased in the order RJ-C<sub>12</sub>-MY<sub>1</sub> > RJ-C<sub>12</sub>-MY<sub>2</sub> > RJ-C<sub>12</sub>-MY<sub>3</sub> > RJ-C<sub>8</sub>-MY<sub>4</sub>, as predicted by DFT. The detailed molecular characterizations in determining the nature of optoelectronic properties i.e.; ground state molar extinction coefficients, excited state molar extinction coefficients, fluorescence quantum yields and excited state lifetimes of newly synthesized Zn(II)porphyrin molecules were examined in order to show the positive correlation between vibrational energy levels, molar extinction coefficients, phonon emitting states and PA effect. Large PA coefficients were observed for the RJ-C<sub>12</sub>-MY<sub>1</sub>, providing an up to ~7-fold PA enhancement over that of ZnTPP used as a reference, which is superior to that of the existing small molecules. Besides offering a superior molecule for PA tomography, the present thesis and criterion adopted here would enable to design simpler molecules with superior photoacoustic and other nonlinear optical properties.



## **TABLE OF CONTENT**

**DECLARATION**

**TITLE PAGE**

**DEDICATION** **ii**

**ACKNOWLEDGEMENTS** **iii**

**ABSTRAK** **iv**

**ABSTRACT** **v**

**TABLE OF CONTENT** **vi**

**LIST OF TABLES** **xi**

**LIST OF FIGURES** **xii**

**LIST OF SYMBOLS** **xviii**

**LIST OF ABBREVIATIONS** **xxi**

**CHAPTER 1 INTRODUCTION** **1**

1.1 Problem Statement 2

1.2 Hypothesis 3

1.3 Research Objectives 4

1.4 Research Scope 5

1.5 Novelty of Research 7

1.6 Thesis Outline 7

**CHAPTER 2 LITERATURE REVIEW** **9**

2.1 Introduction 9

2.2 Metalloporphyrins 11

2.3	Classification of Porphyrins	12
2.3.1	In-Plane Metalloporphyrins	13
2.3.2	SAT Metalloporphyrins	16
2.4	Photochemistry	19
2.4.1	Electronic Properties of Porphyrins	22
2.5	Synthetic Methodologies of Porphyrin	31
2.5.1	Pre-Functionalization	32
2.5.2	Post-Functionalization	41
2.5.3	Metalloporphyrins Synthesis	47
2.6	Density Functional Theory Calculations of Porphyrins	49
2.6.1	Ab-initio Quantum Chemical Methods	50
2.6.2	Density Functional Theory Calculations	51
2.6.3	DFT Calculations by Different Functionals	53
2.6.4	DFT Calculations by Different Basis Sets	55
2.6.5	Time Dependent DFT Calculations	57
2.7	Applications of Porphyrins	58
2.7.1	Photoacoustic Imaging:	59
2.7.2	Porphyrin as Photoacoustic Contrast Agent	62
2.8	Conclusions	65
<b>CHAPTER 3 MATERIALS AND METHODS</b>		<b>66</b>
3.1	Introduction	66
3.2	Research Methodology	66
3.3	Materials and Chemicals	68
3.4	Design of Porphyrins by DFT Calculation	69
3.4.1	Molecular Geometry Optimization	69

3.4.2	Molecular Structures and Nomenclatures	71
3.4.3	Frequency and Vibrational Energy Calculations	72
3.4.4	Excitation Energy Calculations	74
3.5	Synthesis of Newly Designed Zn(II)Porphyrins	76
3.5.1	Instrumentation	76
3.5.2	Synthesis of $\pi$ -Conjugated Meso-Ethynyl-Zn(II)Porphyrins	76
3.5.3	Synthesis of Porphyrins RJ-C <sub>8</sub> -MY <sub>4</sub> :	87
3.6	Photoacoustic Measurements	92
3.7	Summary	93

## **CHAPTER 4 RESULTS AND DISCUSSION: SYNTHESIS AND OPTOELECTRONIC PROPERTIES OF $\pi$ -CONJUGATED MESO-ETHYNYL-ZN(II)PORPHYRINS**

**94**

4.1	Introduction	94
4.2	Synthesis of Meso-Ethynyl-Zn(II)porphyrins	94
4.2.1	Knoevenagel Condensation Reaction	99
4.2.2	Sonogashira Cross-Coupling Reaction	105
4.3	Steady State Absorption Spectral Studies	113
4.3.1	Absorption Spectral Studies Relates to PAI Signals	125
4.4	Steady State Emission Spectral Studies	131
4.4.1	Quantum Yield Studies	143
4.4.2	Time Resolved Photoluminescence Studies	148
4.4.3	PAI Signals Correlation with Emitting Excited States	149
4.5	IR-Vibrational Modes Studies	151
4.5.1	Phonon Emission Correlation with Extended-Conjugation	155
4.6	Thermogravimetric Thermal Analysis (TGA)	156
4.7	Conclusions	157

<b>CHAPTER 5 RESULTS AND DISCUSSION: PHOTOACOUSTIC STUDIES OF <math>\pi</math>-CONJUGATED MESO-ETHYNYL-ZN(II)PORPHYRINS</b>	<b>159</b>
5.1 Introduction	159
5.2 Photoacoustic Studies	159
5.3 Optical and Photoacoustic Z-Scan Studies	160
5.3.1 Dye Concentration Dependence – Low Laser Fluence	162
5.3.2 Dye Concentration Dependence – High Laser Fluence	163
5.4 Photoacoustic Tomography	166
5.5 Optical Photoacoustic Z-Scan Arrangement Details	168
5.6 Photoacoustic Tomography Experimental Details	169
5.7 Conclusions	170
<b>CHAPTER 6 CONCLUSIONS AND RECOMMENDATIONS</b>	<b>171</b>
6.1 Conclusions	171
6.2 Future Recommendations	175
<b>REFERENCES</b>	<b>177</b>
<b>APPENDIX A PORPHYRINS IN NATURE</b>	<b>216</b>
<b>APPENDIX B PORPHYRINS AS PA CONTRAST AGENTS</b>	<b>224</b>
<b>APPENDIX C SCREEN SHORT OF OPTIMIZATION OF PORPHYRINS</b>	<b>232</b>
<b>APPENDIX D MOLECULAR STRUCTURES AND NOMENCLATURES OF DESIGNED PORPHYRINS</b>	<b>233</b>
<b>APPENDIX E UV-VIS SPECTRA OF ALL DESIGNED PORPHYRINS</b>	<b>245</b>
<b>APPENDIX F NMR OF THE SYNTHESIZED PORPHYRINS</b>	<b>248</b>
<b>APPENDIX G ENERGY-GAP LAW</b>	<b>292</b>
<b>APPENDIX H VISUALS OF OPTIMIZED STRUCTURES AND ALL VIBRATIONS OF RJ-C<sub>n</sub>-MY<sub>m</sub> COMPLEXES</b>	<b>294</b>

<b>APPENDIX I MOLECULAR ORBITAL DIAGRAMS OF THE RJ-C<sub>n</sub>-MY<sub>m</sub> SERIES</b>	<b>295</b>
<b>LIST OF PUBLICATIONS</b>	<b>297</b>

## LIST OF TABLES

Table 3.1	Approaches used to optimize the porphyrin molecules	71
Table 4.1	Starting materials and the yields of porphyrin precursors and the meso-ethynyl-Zn(II)porphyrins	96
Table 4.2	Continued	97
Table 4.3	The absorbance spectral data for free-base and Zn(II)porphyrins	118
Table 4.4	Experimental and calculated absorption bands ( $\lambda$ ) alone with their molar extinction coefficients ( $\epsilon$ ) for RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes	130
Table 4.5	The fluorescence spectral data for first series porphyrin complexes	135
Table 4.6	The fluorescence spectral data for second series porphyrin complexes	136
Table 4.7	The fluorescence spectral data for RJ-C <sub>n</sub> -MY <sub>m</sub> series porphyrin complexes	140
Table 4.8	The quantum yield data for RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes	146
Table 5.1	Molar absorption coefficients ( $\epsilon$ ) of RJ-C <sub>n</sub> -MY <sub>m</sub> series of porphyrins at 532 nm & 680 nm	164

## LIST OF FIGURES

Figure 1.1	Basic Structure of Porphyrin	1
Figure 1.2	Excited energy state of a molecule	4
Figure 1.3	The flow chart of research progress and a relationship between the chapters	8
Figure 2.1	Schematic representation of the two tautomers of porphyrin. 1) <i>cis</i> -NH tautomer and 2) <i>trans</i> -NH tautomer	9
Figure 2.2	Metalloporphyrins (adpoted from Gaussview); blue, light gray, dark gray and white colored atoms represents nitrogens, carbons, metal ion and hydrogens	11
Figure 2.3	The most common substitution patterns found in porphyrin macrocycle. 1) porphine; 2) $\beta$ -substituted porphyrin; 3) meso-substituted porphyrin; 4) $\beta$ - and meso-substituted (mixed) porphyrins	13
Figure 2.4	Representation of a) In-plane metalloporphyrins; b) Out-of-plane metalloporphyrin	14
Figure 2.5	Structure representation of In-plane metalloporphyrins; a) Top-view; b) Side-view	14
Figure 2.6	In-plane metalloporphyrin distortion; a) Saddle metalloporphyrin; b) Ruffled metalloporphyrin	15
Figure 2.7	Structure representation of Out-of-plane metalloporphyrin; a) Top-view; b) Side-view	17
Figure 2.8	SAT metalloporphyrin distortion (Dome distortion)	17
Figure 2.9	Formation of In-plane metalloporphyrin via Out-of-plan metallopor-phyrin as intermediate	18
Figure 2.10	Aromaticity in porphyrin: 18- $\pi$ electrons out of 22- $\pi$ electrons form the conjugation pathway (indicated in bold)	19
Figure 2.11	$^1\text{H}$ -NMR representation of porphyrin	20
Figure 2.12	Jablonski diagram showing the decay phenomena for singlet and triplet relaxation after absorption of photon. The solid lines represent the radiation decay and dashed lines for non-radiation decay	21
Figure 2.13	UV-Vis spectra for free-base porphyrins ( $D_{2h}$ ) (top) and metalloporphyrin ( $D_{4h}$ ) (bottom)	24
Figure 2.14	Q-bands absorption spectra for free-base porphyrins: a) etio-type; b) rhodo-type; c) oxo-rhodo-type and d) phyllo-types	24
Figure 2.15	Molecular orbital diagram of “regular” porphyrins	25
Figure 2.16	Molecular orbital diagram of hypso-porphyrins (irregular porphyrins). The notations $a_{1u}$ and $a_{2u}$ are singly degenerate and $e_g$ is doubly degenerate orbitals	26

Figure 2.17	Molecular orbital diagram of hyper-porphyrins (irregular porphyrins)	26
Figure 2.18	Gouterman four-orbital model for $D_{2h}$ symmetry (top) and $D_{4h}$ symmetry porphyrins (bottom)	28
Figure 2.19	The HOMOs and LUMOs orbitals according to the Gouterman Four-Orbital Model	28
Figure 2.20	Emission spectra of metalloporphyrin, regular porphyrin represented in red (ZnTPP) and irregular porphyrin in purple (PtTPP)	29
Figure 2.21	Classification of porphyrin synthesis	31
Figure 2.22	General representation of pre-functionalization of porphyrins	32
Figure 2.23	General methodologies for porphyrin synthesis through tetramerization of mono-pyrrole; i) acid catalyzed condensation; ii) oxidation reaction	33
Figure 2.24	Methodologies for meso-porphyrin synthesis; i) Rothmund synthesis; ii) Adler and Longo synthesis; iii) Lindsey synthesis, acid catalyzed condensation; iiib) Lindsey synthesis, oxidation reaction	34
Figure 2.25	Methodologies for synthesis of unsymmetrical meso-porphyrin; i) acid catalyzed condensation; ii) oxidation reaction	35
Figure 2.26	Methodologies for $\beta$ - porphyrin synthesis	36
Figure 2.27	Self-condensation of 2- substituted pyrrole for $\beta$ - & meso-porphyrin synthesis	37
Figure 2.28	MacDonald (2+2) condensation for porphyrin synthesis	38
Figure 2.29	MacDonald (2+2) condensation for <i>trans</i> substituted meso-porphyrin synthesis	39
Figure 2.30	Tripyrrolic [3+1] route for $\beta$ -unsymmetrical porphyrin synthesis	40
Figure 2.31	Cyclization of linear tetrapyrroles i) MeOH, HCl, RT; ii) MeOH, Cu(OAc) <sub>2</sub> , reflux, 48 h; iii) H <sub>2</sub> SO <sub>4</sub>	40
Figure 2.32	General representation of post-functionalization of porphyrins	41
Figure 2.33	Classification of post-functionalization by position	41
Figure 2.34	Various reaction types of post-functionalization of porphyrins	42
Figure 2.35	Post-functionalization for porphyrins synthesis	43
Figure 2.36	palladium-catalyzed cross-coupling reactions for porphyrins synthesis	45
Figure 2.37	Routes for inserting metal ions into the porphyrin ring through; a) coordination of metal; b) Reduction of metal; c) Oxidation of metal	48
Figure 2.38	Application domains of natural and synthetic porphyrins	59



Figure 2.39	Structures of zinc phthalocyanine, protoporphyrin IX employed as PAI contrast agents	63
Figure 2.40	Structures of uranium porphyrinoid complex employed PAI contrast agents	63
Figure 2.41	Structures of quinoline-annulated porphyrins and bacteriochlorins employed as PAI contrast agents	64
Figure 2.42	Structures of quinoline-annulated porphyrin derivatives employed as PAI contrast agents	64
Figure 3.1	The flow chart of general research methodology	67
Figure 3.2	Geometry optimization steps for the global minimum energies a) compound RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) compound RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) compound RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) compound RJ-C <sub>8</sub> -MY <sub>4</sub>	70
Figure 3.3	Structures and IUPAC naming of porphyrin molecules. The left panel shows planer ChemDraw sketch and right panel are the computed geometry developed using Gauss View. a) compound RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) compound RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) compound RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) compound RJ-C <sub>8</sub> -MY <sub>4</sub>	72
Figure 3.4	IR spectra for a) compound RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) compound RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) compound RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) compound RJ-C <sub>8</sub> -MY <sub>4</sub>	74
Figure 3.5	UV-Vis spectra for a) compound RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) compound RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) compound RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) compound RJ-C <sub>8</sub> -MY <sub>4</sub>	75
Figure 3.6	Synthetic schemes for a) compound RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) compound RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) compound RJ-C <sub>12</sub> -MY <sub>3</sub>	78
Figure 3.7	Synthesis of RJ-C <sub>8</sub> -MY <sub>4</sub>	87
Figure 3.8	Schematic of experimental setup used to obtain nonlinear optical transmission and scattering, and photoacoustic data	93
Figure 4.1	Synthesis of meso-ethynyl-Zn(II)porphyrins	95
Figure 4.2	Reaction mechanism for 1,3-di(alkoxy)benzene (5-5a) and 2,6-bis(alkoxy)benzaldehyde (6-6a)	98
Figure 4.3	General mechanism of Knoevenagel condensation reaction	99
Figure 4.4	Reaction mechanism for 2-cyano-3-(4-iodophenyl)acrylic acid (3)	100
Figure 4.5	Reaction mechanism for di(1H-pyrrol-2-yl)methane (9)	101
Figure 4.6	Reaction mechanism for Free-base porphyrins (10-10b)	102
Figure 4.7	Reaction mechanism for Free-base bromo-porphyrins (11-11b)	104
Figure 4.8	Reaction mechanism for Zn(II) porphyrins (12-12b)	105
Figure 4.9	General mechanism of Sonogashira cross coupling reaction	106
Figure 4.10	General mechanism of copper-free Sonogashira cross coupling reaction	107
Figure 4.11	Reaction mechanism for Zn(II) porphyrins (13-13b)	108

Figure 4.12	Reaction mechanism for Zn(II) porphyrins (14-14b).	109
Figure 4.13	Reaction mechanism for RJ-C <sub>12</sub> -MY <sub>1</sub>	110
Figure 4.14	Reaction mechanism for RJ-C <sub>12</sub> -MY <sub>2</sub> , RJ-C <sub>12</sub> -MY <sub>3</sub> and RJ-C <sub>8</sub> -MY <sub>4</sub>	113
Figure 4.15	Absorption spectra of a) free-base porphyrin (10); b) bromo free-base porphyrin (11); c) bromo Zn(II)porphyrin (12); d) ethynyl-Zn(II)porphyrin (13)	115
Figure 4.16	Q-band absorption spectra of a) free-base porphyrin (10); b) bromo free-base porphyrin (11); c) bromo Zn(II)porphyrin (12); d) ethynyl-Zn(II)porphyrin (13)	116
Figure 4.17	Comparison absorption spectra of free-base porphyrin (10), bromo free-base porphyrin (11), bromo Zn(II)porphyrin (12) and ethynyl-Zn(II)porphyrin (13)	117
Figure 4.18	Absorption spectra of a) free-base porphyrin (10b); b) bromo free-base porphyrin (11b); c) bromo Zn(II)porphyrin (12b); d) ethynyl-Zn(II)porphyrin (13b)	120
Figure 4.19	Q-band absorption spectra of a) free-base porphyrin (10b); b) bromo free-base porphyrin (11b); c) bromo Zn(II)porphyrin (12b); d) ethynyl-Zn(II)porphyrin (13b)	121
Figure 4.20	Comparison absorption spectra of free-base porphyrin (10b), bromo free-base porphyrin (11b), bromo Zn(II)porphyrin (12b) and ethynyl-Zn(II)porphyrin (13b)	121
Figure 4.21	Comparison of absorption spectra between first & second series of porphyrins	122
Figure 4.22	Absorption spectra of RJ-C <sub>12</sub> -MY <sub>1</sub> , RJ-C <sub>12</sub> -MY <sub>2</sub> , RJ-C <sub>12</sub> -MY <sub>3</sub> and RJ-C <sub>8</sub> -MY <sub>4</sub>	123
Figure 4.23	Comparison between absorption spectra of RJ-C <sub>12</sub> -MY <sub>1</sub> , RJ-C <sub>12</sub> -MY <sub>2</sub> , RJ-C <sub>12</sub> -MY <sub>3</sub> and RJ-C <sub>8</sub> -MY <sub>4</sub>	124
Figure 4.24	Comparison between absorption spectra of a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> and d) RJ-C <sub>8</sub> -MY <sub>4</sub> with their respective parent compounds	125
Figure 4.25	Comparison between experimental and calculated absorption spectra of a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> and d) RJ-C <sub>8</sub> -MY <sub>4</sub>	128
Figure 4.26	LUMO orbitals for a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> and d) RJ-C <sub>8</sub> -MY <sub>4</sub>	129
Figure 4.27	Emission spectra of a) free-base porphyrin (10); b) bromo free-base porphyrin (11); c) bromo Zn(II)porphyrin (12); d) ethynyl-Zn(II)porphyrin (13)	133
Figure 4.28	Comparison between absorption and emission spectra of a) free-base porphyrin (10); b) bromo free-base porphyrin (11); c) bromo Zn(II)porphyrin (12); and d) ethynyl-Zn(II)porphyrin (13)	134

Figure 4.29	Comparison of emission spectra of free-base porphyrin (10), bromo free-base porphyrin (11), bromo Zn(II)porphyrin (12) and ethynyl-Zn(II)porphyrin (13)	135
Figure 4.30	Emission spectra of a) free-base porphyrin (10b); b) bromo free-base porphyrin (11b); c) bromo Zn(II)porphyrin (12b); d) ethynyl-Zn(II)porphyrin (13b)	137
Figure 4.31	Comparison absorption and emission spectra of a) free-base porphyrin (10b); b) bromo free-base porphyrin (11b); c) bromo Zn(II)porphyrin (12b); and d) ethynyl-Zn(II)porphyrin (13b)	138
Figure 4.32	Comparison of emission spectra of free-base porphyrin (10b), bromo free-base porphyrin (11b), bromo Zn(II)porphyrin (12b) and ethynyl-Zn(II)porphyrin (13b)	138
Figure 4.33	Emission spectra of RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	139
Figure 4.34	Comparison between absorption and emission spectra of RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	141
Figure 4.35	PL spectra of RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes excited by femtosecond laser at 400 nm wavelength and having 1.4 mW power	142
Figure 4.36	Emission spectra of RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes at different excitation wavelength a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	143
Figure 4.37	Absorbance profile for $\Phi_f$ calculation of RJ-C <sub>n</sub> -MY <sub>m</sub> series of porphyrin complexes at different concentrations a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	144
Figure 4.38	Emission profile for $\Phi_f$ calculation of RJ-C <sub>n</sub> -MY <sub>m</sub> series of porphyrin complexes at different concentrations a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	145
Figure 4.39	Slopes of the line generated from plotting the absorbance against the integrated emission intensity at various wavelengths for RJ-C <sub>12</sub> -MY <sub>1</sub> complex	146
Figure 4.40	Slopes of the line generated from plotting the absorbance against the integrated emission intensity at various wavelengths for RJ-C <sub>12</sub> -MY <sub>2</sub> complex	147
Figure 4.41	Slopes of the line generated from plotting the absorbance against the integrated emission intensity at various wavelengths for RJ-C <sub>12</sub> -MY <sub>3</sub> complex	147
Figure 4.42	Slopes of the line generated from plotting the absorbance against the integrated emission intensity at various wavelengths for RJ-C <sub>8</sub> -MY <sub>4</sub> complex	148
Figure 4.43	a) Time-resolved photoluminescence spectra (TRPL) of RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes b) the fast and slow time components for RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes	149

Figure 4.44	The contribution from vibrational states in the absorption and emission spectra of a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	151
Figure 4.45	Experimental FTIR spectra for RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes; a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	152
Figure 4.46	Computed FTIR spectra for RJ-C <sub>n</sub> -MY <sub>m</sub> series complexes; a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> ; d) RJ-C <sub>8</sub> -MY <sub>4</sub>	153
Figure 4.47	Vibrational spectra of a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> and d) RJ-C <sub>8</sub> -MY <sub>4</sub>	153
Figure 4.48	Comparision between vibrational modes of RJ-C <sub>12</sub> -MY <sub>1</sub> , RJ-C <sub>12</sub> -MY <sub>2</sub> , RJ-C <sub>12</sub> -MY <sub>3</sub> and RJ-C <sub>8</sub> -MY <sub>4</sub>	154
Figure 4.49	TGA curve of a) RJ-C <sub>12</sub> -MY <sub>1</sub> ; b) RJ-C <sub>12</sub> -MY <sub>2</sub> ; c) RJ-C <sub>12</sub> -MY <sub>3</sub> and d) RJ-C <sub>8</sub> -MY <sub>4</sub>	156
Figure 5.1	Optical absorption as a function of laser fluence for the RJ-C <sub>n</sub> -MY <sub>m</sub> series of porphyrins absorbing 0.1 OD in a 2 mm quartz cuvette at 532 nm	161
Figure 5.2	Photoacoustic response plotted with laser fluence. All the porphyrin samples were recorded at 0.1 OD in a 2 mm quartz cuvette at 532 nm	161
Figure 5.3	Optical Z-scan absorption (top) and photoacoustic emission (bottom) of all dyes as a function of optical density and concentration, respectively, measured at 20 mJ/cm <sup>2</sup> fluence	163
Figure 5.4	Plots of optical Z-scan absorption and PA emission vs concentration recorded 400 mJ/cm <sup>2</sup> laser fluence	164
Figure 5.5	Photoacoustic response plotted with laser fluence. All the porphyrin samples were recorded at 0.3 OD in a 2 mm quartz cuvette at 680 nm	165
Figure 5.6	Photoacoustic emission of all the RJ-C <sub>n</sub> -MY <sub>m</sub> series of porphyrin as a function of concentration, respectively, measured at 20 mJ/cm <sup>2</sup> fluence	166
Figure 5.7	PAT image recorded at the low laser fluence of 20 mJ/cm <sup>2</sup> and high laser fluence of 400 mJ/cm <sup>2</sup> at $\lambda_{exc}$ = 532 nm. All the RJ-C <sub>n</sub> -MY <sub>m</sub> samples had 0.1 O.D. at 532 nm in the 1 mm path length borosilicate capillary tubes. (top) Reconstructed WORD FOR WORD image at low laser fluence, normalized with the maximum PA signal strength of high laser fluence. The line plot is the PA signal strength for each sample. (bottom) Reconstructed PAT image obtained at high laser fluence and the corresponding line profile	167
Figure 5.8	PAT image recorded at $\lambda_{exc}$ = 680 nm. All the RJ-C <sub>n</sub> -MY <sub>m</sub> samples had 0.1 O.D. at 680 nm in the 1 mm path length borosilicate capillary tubes	168

## LIST OF SYMBOLS

$\alpha$	Alpha
$\beta$	Beta
cm	Centimetre
$\Delta$	Chemical shift
$C_e$	Concentration of excited state molecules
P	Density
R	Electron position
eV	Electron volt
$\lambda_{em}$	Emission wavelength
$E_g$	Energy-gap
$\lambda_{ex}$	Excitation wavelength
Fs	Femtosecond
$\epsilon_e$	First excited state molar extinction coefficient
$\Phi_f$	Fluorescence quantum yields
$C_g$	Ground state concentration
$\epsilon_g$	Ground state molar extinction coefficient
$\Gamma$	Grüneisen coefficient
$t_{1/2}$	Half-life
Hz	Hertz
$I$	Incident photon fluence
$\nabla^2$	Laplacian operator
MHz	Megahertz
$\mu J$	Microjoules

$\mu\text{m}$	Micrometre
mJ	Millijoules
mL	Millilitre
Mm	Millimeter
mmol	Millimole
Ms	Millisecond
M	Molar
E	Molar extinction coefficient
Nm	Nanometre
Ns	Nanosecond
$k_{\text{nr}}$	Non-radiative decay
F	Oscillation strengths
Ppm	Parts per million
$\Phi_{\text{ph}}$	Phosphorescence quantum yield
$\pi$	Pi
Pm	Picometer
Ps	Picosecond
$\hbar$	Plank constant
$\Phi_{\text{nr}}$	Quantum yield for non-radiative decay
$k_{\text{r}}$	Radiative decay
N	Refractive index
$^1\tau$	Singlet excited state life time
$C_{\text{p}}$	Specific heat of the medium at constant pressure
$C_{\text{v}}$	Specific heat of the medium at constant volume

$C$	Speed of light
$\Delta\nu_1$	Splitting between excited states Q(1,0) and Q(0,0) vibrational bands
$\Delta\nu_2$	Splitting between ground states Q(0,0) and Q(1,0) vibrational bands
$\Delta\nu$	Stokes shift
$\alpha$	Thermal expansion coefficient/ Linear absorption coefficient
$V_s$	Velocity of sound
$W$	Watt
$\Psi$	Wave function
$\lambda$	Wavelength

## LIST OF ABBREVIATIONS

B3LYP	Becke, 3-parameter, Lee-Yang-Parr
(Pd(PPh <sub>3</sub> ) <sub>2</sub> Cl <sub>2</sub> )	Bis(triphenylphosphine)palladium(II)chloride
BODIPY	Boron-Dipyrromethene
<sup>13</sup> CNMR	Carbon Nuclear Magnetic Resonance
CHCl <sub>3</sub>	Chloroform
CI	Configuration Interaction
CuI	Copper iodide
CAMB3LYP	Coulomb attenuated-B3LYP
CC	Coupled-Cluster
CCSD	Coupled Cluster with single and double excitations
DFT	Density Functional Theory
CDCl <sub>3</sub>	Deuteriochloroform
CD <sub>3</sub> OH	Deuteromethanol
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
CH <sub>2</sub> Cl <sub>2</sub> or DCM	Dichloromethane
DMF	Dimethylformamide
DZ	Double Zeta
E <sub>em</sub>	Emission energy
EtOAc	Ethyl acetate
E <sub>ex</sub>	Excitation energy
S <sub>1</sub>	Excited singlet state
FDA	Food and Drug Administration
FTIR	Fourier Transform Infrared
GTOs	Gaussian type orbitals
GUI	Graphical user interface



S <sub>0</sub>	Ground state
HF	Hartree-Fock
HOMO	Highest Occupied Molecular Orbital
HRMS	High Resonance Mass Spectroscopy
HK	Hohenberg-Kohn
HI	Hydriodic acid
HCl	Hydrochloric acid
HBr	Hydrogen bromide
H <sub>2</sub> O <sub>2</sub>	Hydrogen peroxide
ICG	Indocyanine green
IR	Infrared
IEF-PCM	Integral Equation Formalism Polarizable Continuum Model
IC	Interconversion
IUPAC	International union for pure and applied chemistry
ISC	Intersystem crossing
LR-TD-DFT	Linear response Time-dependent Density Functional Theory
LANL2DZ	Los Alamos National Laboratory (LANL) ECP with Double Zeta valence
LUMO	Lowest Unoccupied Molecular Orbital
MgSO <sub>4</sub>	Magnesium sulfate
MATLAB	MATrix LABoratory
H <sub>2</sub> TMP	meso-tetramethylporphyrin
H <sub>2</sub> TPP	meso-tetraphenylporphyrin
M <sup>II</sup> (L)(CO)	Metal ligand cobalt
MX <sub>2</sub>	Metal salts
MLCT	Metal to ligand charge transfer
CH <sub>3</sub> OH or MeOH	Methanol

MP2	Møller-Plesset Perturbation
NBS	<i>N</i> -bromosuccinimide
nBuLi	<i>n</i> -Butyllithium
NCS	<i>N</i> -chlorosuccinimide
NIR	Near-Infrared
Nd:YAG	Neodymium Yttrium Aluminum Garnet
NIS	<i>N</i> -iodosuccinimide
HNO <sub>3</sub>	Nitric acid
N <sub>2</sub>	Nitrogen gas
N <sub>2</sub> O <sub>4</sub>	Nitrogen tetroxide
NO <sub>2</sub> BF <sub>4</sub>	Nitrosonium borofluorate
NMR	Nuclear Magnetic Resonance
OPAZ	Optical and Photoacoustic Z-Scan
OD	Optical density
PhSeBr <sub>3</sub>	Phenylselenium tribromide
PhSeCl <sub>3</sub>	Phenylselenium trichloride
POCl <sub>3</sub>	Phosphoryl chloride
PA	Photoacoustic
PAI	Photoacoustic Imaging
PAM	Photoacoustic Microscopy
PAT	Photoacoustic Tomography
PCM	Polarizable Continuum Model
K <sub>2</sub> CO <sub>3</sub>	Potassium carbonate
<sup>1</sup> HNMR	Proton Nuclear Magnetic Resonance
QZ	Quadruple Zeta
QC	Quantum chemical

RSA	Reverse saturable absorption
SCRf	Self-consistent reaction-field
SAT	Sitting at top
STOs	Slater type orbital
NaN <sub>3</sub>	Sodium azide
NaBH <sub>4</sub>	Sodium borohydride
NaOH	Sodium hydroxide
TBAF	Tetrabutylammonium fluoride
THF	Tetrahydrofuran
TMEDA	Tetramethylethylenediamine
TGA	Thermal Gravimetric Analysis
TLC	Thin-Layer Chromatograph
TD-DFT	Time-dependent Density Functional Theory
TRPL	Time-resolved Photoluminescence
SnCl <sub>2</sub>	Tin(II) chloride
NEt <sub>3</sub>	Triethylamine
TFA	Trifluoroacetic acid
TIPS	Triisopropylsilyl
AsPh <sub>3</sub>	Triphenylarsine
T <sub>1</sub>	Triplet excited state
T <sub>n</sub>	Triplet excited states
TZ	Triple Zeta
Pd <sub>2</sub> (dba) <sub>3</sub>	Tris(dibenzylideneacetone)dipalladium(0)
UV-Vis	UltraViolet-Visible
VR	Vibrational relaxation
Zn(OAc) <sub>2</sub> •2H <sub>2</sub> O	Zinc acetate dehydrate
ZnTPP	Zinc tetraphenylporphyrin

## REFERENCES

- Abedian, N. and Dehghani, H. (2013). Novel molecular complexation between meso-tetraarylporphyrinato magnesium(II) and phosphorus(III) chloride. *Inorganic Chemistry Communications*, 36, 77-80.
- Abraham, R. J., Hawkes, G. E., Hudson, M. F. and Smith, K. M. (1975). The nuclear magnetic resonance spectra of porphyrins. Part X. Carbon-13 nuclear magnetic resonance spectra of some meso-tetraarylporphyrins and their metal chelates. *Journal of the Chemical Society, Perkin Transactions*, 2(3), 204-211.
- Abrahamse, H. and Hamblin, M. R. (2016). New photosensitizers for photodynamic therapy. *Biochemical Journal*, 473(4), 347-364.
- Abuteen, A., Zanganeh, S., Akhigbe, J., Samankumara, L. P., Aguirre, A., Biswal, N. . . Brückner, C. (2013). The evaluation of NIR-absorbing porphyrin derivatives as contrast agents in photoacoustic imaging. *Physical Chemistry Chemical Physics*, 15(42), 18502-18509.
- Adar, F. (1978). Electronic absorption spectra of hemes and hemoproteins. In D. Dolphin (Ed.), *The Porphyrins* (Vol 3, pp 167-209). New York: Academic Press.
- Adler, A. D. and Gouterman, M. (1985). *An historical survey of porphyrins, their related spectra and their interpretation*. Paper presented at the Optical properties and structure of tetrapyrroles: proceedings of a symposium held at the University of Konstanz, West Germany, August 12-17, 1984.
- Adler, A. D., Longo, F. R., Finarelli, J. D., Goldmacher, J., Assour, J. and Korsakoff, L. (1967). A simplified synthesis for meso-tetraphenylporphine. *The Journal of Organic Chemistry*, 32(2), 476-476.
- Adler, A. D., Longo, F. R., Kampas, F. and Kim, J. (1970). On the preparation of metalloporphyrins. *Journal of Inorganic and Nuclear Chemistry*, 32(7), 2443-2445.
- Aihara, H., Jaquinod, L., Nurco, D. J. and Smith, K. M. (2001). Multicarbocycle Formation Mediated by Arenoporphyrin 1, 4-Diradicals: Synthesis of Picenoporphyrins. *Angewandte Chemie International Edition*, 40(18), 3439-3441.
- Aizawa, K., Okunaka, T., Ohtani, T., Kawabe, H., Yasunaka, Y., O'Hata, S. . . Kato, H. (1987). Localization of mono-l-aspartyl chlorin e6 (npe6) in mouse tissues. *Photochemistry and Photobiology*, 46(5), 789-793.

- Akins, D., Zhu, H.-R. and Guo, C. (1994). Absorption and Raman scattering by aggregated meso-tetrakis (p-sulfonatophenyl) porphine. *The Journal of Physical Chemistry*, 98(14), 3612-3618.
- Akins, D. L., Özçelik, S., Zhu, H.-R. and Guo, C. (1996). Fluorescence decay kinetics and structure of aggregated tetrakis (p-sulfonatophenyl) porphyrin. *The Journal of Physical Chemistry*, 100(34), 14390-14396.
- Al-Karadaghi, S., Franco, R., Hansson, M., Shelnutt, J. A., Isaya, G. and Ferreira, G. C. (2006). Chelataases: distort to select? *Trends in Biochemical Sciences*, 31(3), 135-142.
- Ali, H. and van Lier, J. E. (1991). Phenylselenenyl halides: Efficient reagents for the selective halogenation and nitration of porphyrins. *Tetrahedron Letters*, 32(38), 5015-5018.
- Ali, H. and van Lier, J. E. (1994). Synthesis of  $\beta$ -substituted porphyrins using palladium catalysed reactions. *Tetrahedron*, 50(41), 11933-11944.
- Amao, Y., Asai, K., Miyakawa, K. and Okura, I. (2000). Oxygen sensing using palladium porphyrin with long alkyl chain self-assembled film. *Journal of Porphyrins and Phthalocyanines*, 4(1), 19-22.
- Ambroise, A., Wagner, R. W., Rao, P. D., Riggs, J. A., Hascoat, P., Diers, J. R. . . . Holten, D. (2001). Design and synthesis of porphyrin-based optoelectronic gates. *Chemistry of Materials*, 13(3), 1023-1034.
- Anderson, H. L. (1994). Conjugated porphyrin ladders. *Inorganic Chemistry*, 33(5), 972-981.
- Andrews, L. E., Bonnett, R., Kozyrev, A. N. and Appelman, E. H. (1988). meso-Reactivity of porphyrins and related compounds. Part 10. Direct fluorination of octaethylporphyrin with caesium fluoroxydisulfate. *Journal of the Chemical Society, Perkin Transactions*, 1(7), 1735-1738.
- Andzelm, J., Rinderspacher, B. C., Rawlett, A., Dougherty, J., Baer, R. and Govind, N. (2009). Performance of DFT methods in the calculation of optical spectra of TCF-chromophores. *Journal of Chemical Theory and Computation*, 5(10), 2835-2846.
- Aoyagi, K., Haga, T., Toi, H., Aoyama, Y., Mizutani, T. and Ogoshi, H. (1997). Electron Deficient Porphyrins III. Facile Syntheses of Perfluoroalkylporphyrins Including Water Soluble Porphyrin. *Bulletin of the Chemical Society of Japan*, 70(4), 937-943.
- Aratani, N., Kim, D. and Osuka, A. (2009).  $\pi$ -Conjugation Enlargement Toward the Creation of Multi-Porphyrinic Systems with Large Two-Photon Absorption Properties. *Chemistry—An Asian Journal*, 4(8), 1172-1182.

- Atefi, F. and Arnold, D. P. (2008). Porphyrins with metal, metalloid or phosphorus atoms directly bonded to the carbon periphery. *Journal of Porphyrins and Phthalocyanines*, 12(07), 801-831.
- Atkinson, S., Brady, S., James, J. and Nolan, K. (1995). Synthetic haems as mimics for high valent intermediates in haemoprotein catalysed oxidations. Synthesis and oxidation of chloro-7, 8, 17, 18-tetracyano-5, 10, 15, 20-tetraphenylporphyrinatoiron (III), a haem which contains strongly electron-withdrawing groups in the  $\beta$ -pyrrole positions. *Pure and Applied Chemistry*, 67(7), 1109-1116.
- Baba, H., Chen, J., Shinokubo, H. and Osuka, A. (2008). Efficient Rhodium-Catalyzed Installation of Unsaturated Ester Functions onto Porphyrins: Site-Specific Heck-Type Addition versus Conjugate Addition. *Chemistry-A European Journal*, 14(14), 4256-4262.
- Bakar, M. A., Sergeeva, N. N., Juillard, T. and Senge, M. O. (2011). Synthesis of ferrocenyl porphyrins via suzuki coupling and their photophysical properties. *Organometallics*, 30(11), 3225-3228.
- Barkigia, K., Fajer, J., Adler, A. and Williams, G. (1980). Crystal and molecular structure of (5, 10, 15, 20-tetra-N-propylporphinato) lead (II): a "roof" porphyrin. *Inorganic Chemistry*, 19(7), 2057-2061.
- Barkigia, K. M., Berber, M. D., Fajer, J., Medforth, C. J., Renner, M. W. and Smith, K. M. (1990). Nonplanar porphyrins. X-ray structures of (2, 3, 7, 8, 12, 13, 17, 18-octaethyl-and-octamethyl-5, 10, 15, 20-tetraphenylporphinato) zinc (II). *Journal of the American Chemical Society*, 112(24), 8851-8857.
- Barnes, J. and Dorough, G. (1950). Exchange and Replacement Reactions of  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ -Tetraphenyl-metalloporphyrins. *Journal of the American Chemical Society*, 72(9), 4045-4050.
- Barnett, G. H., Hudson, M. F. and Smith, K. M. (1975). Concerning meso-tetraphenylporphyrin purification. *Journal of the Chemical Society, Perkin Transactions*, 1(14), 1401-1403.
- Bartoli, J., Battioni, P., De Foor, W. and Mansuy, D. (1994). Synthesis and remarkable properties of iron  $\beta$ -polynitroporphyrins as catalysts for monooxygenation reactions. *Journal of the Chemical Society. Chemical Communications*, (1), 23-24.
- Baruah, T. and Pederson, M. R. (2009). DFT calculations on charge-transfer states of a Carotenoid-Porphyrin-C60 molecular triad. *Journal of Chemical Theory and Computation*, 5(4), 834-843.

- Battersby, A. R., Hunt, E., Ihara, M., McDonald, E., Paine, J. B., Satoh, F. and Saunders, J. (1974). Structure of a heptacarboxylic porphyrin enzymically derived from porphobilinogen. *Journal of the Chemical Society, Chemical Communications*, (23), 994-995.
- Bauschlicher, C. W. (1995). A comparison of the accuracy of different functionals. *Chemical Physics Letters*, 246(1), 40-44.
- Bayer, C. L., Wlodarczyk, B. J., Finnell, R. H. and Emelianov, S. Y. (2017). Ultrasound-guided spectral photoacoustic imaging of hemoglobin oxygenation during development. *Biomedical Optics Express*, 8(2), 757-763.
- Becke, A. D. (1988). Density-functional exchange-energy approximation with correct asymptotic behavior. *Physical Review A*, 38(6), 3098-3100.
- Becke, A. D. (1992). Density-functional thermochemistry. I. The effect of the exchange-only gradient correction. *The Journal of Chemical Physics*, 96(3), 2155-2160.
- Becke, A. D. (1993). Becke's three parameter hybrid method using the LYP correlation functional. *Journal of Chemical Physics*, 98, 5648-5652.
- Beyzavi, M. H., Lentz, D., Reissig, H. U. and Wiehe, A. (2013). Synthesis of Functionalized, Sterically Congested Calix [4] phyrin Macrocycles Using Donor–Acceptor-Substituted Cyclopropanes—First Example of a Mono-meso-spirolactone Incorporated into a Calix [4] phyrin. *European Journal of Organic Chemistry*, 2013(2), 269-282.
- Bhyrappa, P. and Krishnan, V. (1991). Octabromotetraphenylporphyrin and its metal derivatives: Electronic structure and electrochemical properties. *Inorganic Chemistry*, 30(2), 239-245.
- Bhyrappa, P. and Purushothaman, B. (2001). Perbrominated 2-nitrotetraphenylporphyrins: electrochemical and axial ligation properties. *Journal of the Chemical Society, Perkin Transactions*, 2(2), 238-242.
- Binstead, R. A., Crossley, M. J. and Hush, N. S. (1991). Modulation of valence orbital levels of metalloporphyrins by. beta.-substitution: evidence from spectroscopic and electrochemical studies of 2-substituted metallo-5, 10, 15, 20-tetraphenylporphyrins. *Inorganic Chemistry*, 30(6), 1259-1264.
- Böhm, P. and Gröger, H. (2015). Iron (III)-porphyrin Complex FeTSPP: A Versatile Water-soluble Catalyst for Oxidations in Organic Syntheses, Biorenewables Degradation and Environmental Applications. *ChemCatChem*, 7(1), 22-28.
- Bonnett, R., Gale, I. and Stephenson, G. (1966). The meso-reactivity of porphyrins and related compounds. Part II. Halogenation. *Journal of the Chemical Society C: Organic*, 1600-1604.

- Bonnett, R. and Stephenson, G. (1965). The meso reactivity of porphyrins and related compounds. I. Nitration. *The Journal of Organic Chemistry*, 30(8), 2791-2798.
- Borah, K. D. and Bhuyan, J. (2017). Magnesium porphyrins with relevance to chlorophylls. *Dalton Transactions*, 46(20), 6497-6509.
- Boudif, A. and Momenteau, M. (1994). Synthesis of a porphyrin-2, 3-diacrylic acid using a new '3+ 1' type procedure. *Journal of the Chemical Society, Chemical Communications*, (18), 2069-2070.
- Boudif, A. and Momenteau, M. (1996). A new convergent method for porphyrin synthesis based on a '3+ 1' condensation. *Journal of the Chemical Society, Perkin Transactions, 1*(11), 1235-1242.
- Boyle, R. W., Johnson, C. K. and Dolphin, D. (1995). Iodination and Heck alkynylation of 5, 15-diphenylporphyrin. A convenient entry to asymmetrically meso-substituted porphyrins. *Journal of the Chemical Society, Chemical Communications*, (5), 527-528.
- Boys, S. F. (1950). *Electronic wave functions. I. A general method of calculation for the stationary states of any molecular system*. Paper presented at the Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences.
- Braslavsky, S. E. and Heibel, G. E. (1992). Time-resolved photothermal and photoacoustic methods applied to photoinduced processes in solution. *Chemical Reviews*, 92(6), 1381-1410.
- Broadhurst, M., Grigg, R. and Johnson, A. (1971). Synthesis of porphin analogues containing furan and/or thiophen rings. *Journal of the Chemical Society C: Organic*, 3681-3690.
- Browning, L. M., Lee, K. J., Huang, T., Nallathamby, P. D., Lowman, J. E. and Nancy Xu, X.-H. (2009). Random walk of single gold nanoparticles in zebrafish embryos leading to stochastic toxic effects on embryonic developments. *Nanoscale*, 1(1), 138-152.
- Brückner, C., Hyland, M. A., Sternberg, E. D., MacAlpine, J. K., Rettig, S. J., Patrick, B. O. and Dolphin, D. (2005). Preparation of [meso-tetraphenylchlorophinato] nickel (II) by stepwise deformylation of [meso-tetraphenyl-2, 3-diformyl-secochlorinato] nickel (II): conformational consequences of breaking the structural integrity of nickel porphyrins. *Inorganica Chimica Acta*, 358(10), 2943-2953.
- Buchler, J. (1978). Synthesis and properties of metalloporphyrins. In D. Dolphin (ed.), *The porphyrins* (Vol. 1, pp. 389). New York: Academic Press.



- Burrell, A. K. and Officer, D. L. (1998). Functionalizing porphyrins via Wittig reactions: a building block approach. *Synlett*, 1998(12), 1297-1307.
- Callot, H. (1974). Bromination of meta-tetraphenylporphine-preparation of alkyl derivatives and polycyanoporphines. *Bulletin De La Societe Chimique De France Partie Ii-Chimie Moleculaire Organique Et Biologique* (7-8), 1492-1496.
- Calvin, M., Ball, R. and Aronoff, S. (1943).  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ -Tetraphenylchlorin. *Journal of the American Chemical Society*, 65(11), 2259-2259.
- Caspar, J. V. and Meyer, T. J. (1983). Application of the energy gap law to nonradiative, excited-state decay. *The Journal of Physical Chemistry*, 87(6), 952-957.
- Cavaleiro, J. A., Gonsalves, A. M. d. A. R., Kenner, G. W. and Smith, K. M. (1973). Pyrroles and related compounds. Part XXII. Syntheses of pyrromethanes and a tripyrrane. *Journal of the Chemical Society, Perkin Transactions, 1*, 2471-2478.
- Cavaleiro, J. A., Gonsalves, A. M. d. A. R., Kenner, G. W. and Smith, K. M. (1974). Pyrroles and related compounds. Part XXXIII. Total synthesis of deuteriated derivatives of protoporphyrin-IX for nuclear magnetic resonance studies of haemoproteins. *Journal of the Chemical Society, Perkin Transactions, 1*, 1771-1781.
- Chai, J.-D. and Head-Gordon, M. (2008a). Long-range corrected hybrid density functionals with damped atom-atom dispersion corrections. *Physical Chemistry Chemical Physics*, 10(44), 6615-6620.
- Chai, J.-D. and Head-Gordon, M. (2008b). Systematic optimization of long-range corrected hybrid density functionals. *The Journal of Chemical Physics*, 128(8), 084106 (1-15).
- Chandra, T., Kraft, B. J., Huffman, J. C. and Zaleski, J. M. (2003). Synthesis and structural characterization of porphyrinic enediynes: Geometric and electronic effects on thermal and photochemical reactivity. *Inorganic Chemistry*, 42(17), 5158-5172.
- Chaves, O. A., Jesus, C. S., Henriques, E. S., Brito, R. M. and Serpa, C. (2016). In situ ultra-fast heat deposition does not perturb the structure of serum albumin. *Photochemical & Photobiological Sciences*, 15(12), 1524-1535.
- Check, C. E., Faust, T. O., Bailey, J. M., Wright, B. J., Gilbert, T. M. and Sunderlin, L. S. (2001). Addition of polarization and diffuse functions to the LANL2DZ basis set for p-block elements. *The Journal of Physical Chemistry A*, 105(34), 8111-8116.

- Chen, D.-M., Zhang, Y.-H., He, T.-J. and Liu, F.-C. (2002). Raman and UV–visible absorption spectra of ion-paired aggregates of copper porphyrins. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 58(10), 2291-2297.
- Chen, J., Aratani, N., Shinokubo, H. and Osuka, A. (2009). Post-Modification of meso–meso-Linked Porphyrin Arrays by Iridium and Rhodium Catalyses for Tuning of Energy Gap. *Chemistry–An Asian Journal*, 4(7), 1126-1133.
- Chen, Y.-J., Lee, G.-H., Peng, S.-M. and Yeh, C.-Y. (2005). Unexpected formation of porphyrinic enyne under Sonogashira conditions. *Tetrahedron Letters*, 46(9), 1541-1544.
- Chen, Y.-S., Frey, W., Kim, S., Kruizinga, P., Homan, K. and Emelianov, S. (2011). Silica-coated gold nanorods as photoacoustic signal nanoamplifiers. *Nano Letters*, 11(2), 348-354.
- Chen, Y. and Zhang, X. P. (2003). Facile and Efficient Synthesis of meso-Arylamino- and Alkylamino-Substituted Porphyrins via Palladium-Catalyzed Amination. *The Journal of Organic Chemistry*, 68(11), 4432-4438.
- Chinchilla, R. and Nájera, C. (2007). The Sonogashira reaction: a booming methodology in synthetic organic chemistry. *Chemical Reviews*, 107(3), 874-922.
- Chinchilla, R. and Nájera, C. (2011). Recent advances in Sonogashira reactions. *Chemical Society Reviews*, 40(10), 5084-5121.
- Chizhova, N., Kumeev, R. and Mamardashvili, N. Z. (2013). Synthesis and spectral properties of cobalt (II) and cobalt (III) tetraarylporphyrinates. *Russian Journal of Inorganic Chemistry*, 58(6), 740-743.
- Chorghade, M., Dolphin, D., Dupre, D., Hill, D., Lee, E. and Wijesekera, T. (1996). Improved protocols for the synthesis and halogenation of sterically hindered metalloporphyrins. *Synthesis*, 1996(11), 1320-1324.
- Christodoulides, D. N., Khoo, I. C., Salamo, G. J., Stegeman, G. I. and Van Stryland, E. W. (2010). Nonlinear refraction and absorption: mechanisms and magnitudes. *Advances in Optics and Photonics*, 2(1), 60-200.
- Chumakov, D., Khoroshutin, A., Anisimov, A. and Kobrakov, K. (2009). Bromination of porphyrins. *Chemistry of Heterocyclic Compounds*, 45(3), 259-283.
- Čížek, J. (1966). On the correlation problem in atomic and molecular systems. Calculation of wavefunction components in Ursell-type expansion using quantum-field theoretical methods. *The Journal of Chemical Physics*, 45(11), 4256-4266.

- Collman, J., Barnes, C., Brothers, P., Collins, T., Ozawa, T., Gallucci, J. and Ibers, J. A. (1984). Oxidation of ruthenium (II) and ruthenium (III) porphyrins. Crystal structures of  $\mu$ -oxo-bis [(p-methylphenoxo)(meso-tetraphenylporphyrinato) ruthenium (IV)] and ethoxo (meso-tetraphenylporphyrinato)(ethanol) ruthenium (III)-bisethanol. *Journal of the American Chemical Society*, 106(18), 5151-5163.
- Collman, J. P., Barnes, C. E., Swepston, P. N. and Ibers, J. A. (1984). Synthesis, proton NMR, and structural characterization of binuclear ruthenium porphyrin dimers. *Journal of the American Chemical Society*, 106(12), 3500-3510.
- Cook, L., Brewer, G. and Wong-Ng, W. (2017). Structural aspects of porphyrins for functional materials applications. *Crystals*, 7(7), 223(1-22).
- Cramer, C. J. (2013). *Essentials of computational chemistry: theories and models*. USA: John Wiley & Sons.
- Cunha, A. C., Gomes, A. T., Ferreira, V. F., de Souza, M. C., Neves, M. G., Tomé, A. C. . . . Cavaleiro, J. A. (2010). Synthesis of  $\beta$ -Arylporphyrins and Oligophenylenediporphyrins by the Suzuki-Miyaura Reaction. *Synthesis*, 2010(03), 510-514.
- Curtiss, L. A., Redfern, P. C. and Raghavachari, K. (2005). Assessment of Gaussian-3 and density-functional theories on the G3/05 test set of experimental energies. *The Journal of Chemical Physics*, 123(12), 124107(1-12).
- D'Souza, F. and Ito, O. (2005). Photoinduced electron transfer in supramolecular systems of fullerenes functionalized with ligands capable of binding to zinc porphyrins and zinc phthalocyanines. *Coordination Chemistry Reviews*, 249(13), 1410-1422.
- da GH Vicente, M. and M Smith, K. (2014). Syntheses and functionalizations of porphyrin macrocycles. *Current Organic Synthesis*, 11(1), 3-28.
- Dahal, S. and Krishnan, V. (1995). Charge transfer excited states of zinc (II) derivatives of  $\beta$ -substituted dinitrotetraphenylporphyrin. *Journal of Photochemistry and Photobiology A: Chemistry*, 89(2), 105-112.
- Danquah, M. (2017). Review of Handbook of Porphyrin Science with Application to Chemistry, Physics, Materials Science, Engineering, Biology and Medicine. Volume 40: Nanoorganization of Porphyrinoids. *Journal Of Natural Products*, 80(4), 1232–1232.
- Davidson, E. R. and Feller, D. (1986). Basis set selection for molecular calculations. *Chemical Reviews*, 86(4), 681-696.
- de Almeida, J. B., Kenner, G., Rimmer, J. and Smith, K. (1976). Pyrroles and related compounds—XXXV: A stepwise, general synthesis of unsymmetrically substituted porphyrins. *Tetrahedron*, 32(14), 1793-1799.

- de la Zerda, A., Bodapati, S., Teed, R., May, S. Y., Tabakman, S. M., Liu, Z. . . . Gambhir, S. S. (2012). Family of enhanced photoacoustic imaging agents for high-sensitivity and multiplexing studies in living mice. *ACS Nano*, 6(6), 4694-4701.
- Del Bene, J. E., Person, W. B. and Szczepaniak, K. (1995). Properties of Hydrogen-Bonded Complexes Obtained from the B3LYP Functional with 6-31G (d, p) and 6-31+ G (d, p) Basis Sets: Comparison with MP2/6-31+ G (d, p) Results and Experimental Data. *The Journal of Physical Chemistry*, 99(27), 10705-10707.
- Deshpande, R., Jiang, L., Schmidt, G., Rakovan, J., Wang, X., Wheeler, K. and Wang, H. (2009). A Concise Approach to the Synthesis of opp-Dibenzoporphyrins through the Heck Reaction. *Organic Letters*, 11(19), 4251-4253.
- Devillers, C. H., Hebié, S., Lucas, D., Cattey, H., Clément, S. and Richeter, S. (2014). Aromatic Nucleophilic Substitution (S<sub>N</sub>Ar) of meso-Nitroporphyrin with Azide and Amines as an Alternative Metal Catalyst Free Synthetic Approach To Obtain meso-N-Substituted Porphyrins. *The Journal of Organic Chemistry*, 79(14), 6424-6434.
- Dierksen, M. and Grimme, S. (2004). Density functional calculations of the vibronic structure of electronic absorption spectra. *The Journal of Chemical Physics*, 120(8), 3544-3554.
- DiMagno, S. G., Lin, V. S. and Therien, M. J. (1993). Facile elaboration of porphyrins via metal-mediated cross-coupling. *The Journal of Organic Chemistry*, 58(22), 5983-5993.
- Djelal, B. and áGeorge Truscott, T. (1999). Photophysical properties of 5, 10, 15, 20-tetrakis (m-hydroxyphenyl) porphyrin (m-THPP), 5, 10, 15, 20-tetrakis (m-hydroxyphenyl) chlorin (m-THPC) and 5, 10, 15, 20-tetrakis (m-hydroxyphenyl) bacteriochlorin (m-THPBC): a comparative study. *Journal of the Chemical Society, Perkin Transactions*, 2(2), 325-328.
- Dogutan, D. K., Ptaszek, M. and Lindsey, J. S. (2007). Direct synthesis of magnesium porphine via 1-formyldipyrrromethane. *The Journal of Organic Chemistry*, 72(13), 5008-5011.
- Dogutan, D. K., Ptaszek, M. and Lindsey, J. S. (2008). Rational or statistical routes from 1-acyldipyrrromethanes to meso-substituted porphyrins. Distinct patterns, multiple pyridyl substituents, and amphipathic architectures. *The Journal of Organic Chemistry*, 73(16), 6187-6201.
- Dolphin, D. (Ed.). (1978). *The Porphyrins: Structure and Synthesis, Part A* (Vol. 1). London: Academic Press.

- Dong, H., Hou, T., Zhao, Y., Fu, X. and Li, Y. (2012). DFT study of cobalt porphyrin complex for living radical polymerization of olefins. *Computational and Theoretical Chemistry*, 1001, 51-59.
- Drabkin, D. L. (2012). Selected landmarks in the history of porphyrins and their biologically functional derivatives. In D. Dolphin (ed.), *The porphyrins, Structure and Synthesis, Part 1* (Vol. 1, pp. 29-71) New York: Academic Press.
- Drain, C. M., Kirmaier, C., Medforth, C. J., Nurco, D. J., Smith, K. M. and Holten, D. (1996). Dynamic photophysical properties of conformationally distorted nickel porphyrins. 1. Nickel (II) dodecaphenylporphyrin. *The Journal of Physical Chemistry*, 100(29), 11984-11993.
- Dumoulin, F. and Ahsen, V. (2011). Click chemistry: the emerging role of the azide-alkyne Huisgen dipolar addition in the preparation of substituted tetrapyrrolic derivatives. *Journal of Porphyrins and Phthalocyanines*, 15(07), 481-504.
- Duncan, M. A. (2001). *Advances in Metal and Semiconductor Clusters: Metal Ion Solvation and Metal-Ligand Interactions*. Amsterdam: Elsevier.
- Dwyer, P. N., Puppe, L., Buchler, J. and Scheidt, W. R. (1975). Molecular stereochemistry of (alpha.,. gamma.-dimethyl-. alpha.,. gamma.-dihydrooctaethylporphinato) oxotitanium (IV). *Inorganic Chemistry*, 14(8), 1782-1785.
- Éll, A. H., Csjernyik, G., Slagt, V. F., Bäckvall, J. E., Berner, S., Puglia, C. . . . Oscarsson, S. (2006). Synthesis of Thioacetate-Functionalized Cobalt (II) Porphyrins and Their Immobilization on Gold Surface—Characterization by X-ray Photoelectron Spectroscopy. *European Journal of Organic Chemistry*, 2006(5), 1193-1199.
- Elschenbroich, C. (2006). Organometallics, Third Completely Revised and Extended Edition: Wiley-VCH, Weinheim. *Journal Of the American Chemical Society*, 128(46), 12029–12030.
- Falk, J. E. (1964). *Porphyrins and metalloporphyrins: their general, physical and coordination chemistry, and laboratory methods*. Amsterdam, New York: Elsevier Publishing Company.
- Feng, X., Gao, F., Xu, C., Gaoming, L. and Zheng, Y. (2015). Self temperature regulation of photothermal therapy by laser-shared photoacoustic feedback. *Optics Letters*, 40(19), 4492-4495.
- Filonov, G. S., Krumholz, A., Xia, J., Yao, J., Wang, L. V. and Verkhusha, V. V. (2012). Deep-Tissue Photoacoustic Tomography of a Genetically Encoded Near-Infrared Fluorescent Probe. *Angewandte Chemie International Edition*, 51(6), 1448-1451.

- Fischer, H. and Zeile, K. (1929). Synthese des haematoporphyrins, protoporphyrins und haemins. *European Journal of Organic Chemistry*, 468(1), 98-116.
- Fock, V. (1930). Approximation method for the solution of the quantum mechanical multibody problems. *Zeitschrift fur Physik*, 61, 126-148.
- Forbrich, A., Shao, P., Shi, W. and Zemp, R. J. (2016). Lifetime-weighted photoacoustic imaging. *Journal of Optics*, 18(12), 124001(1- 6).
- Frampton, M. J., Akdas, H., Cowley, A. R., Rogers, J. E., Slagle, J. E., Fleitz, P. A. . . . Anderson, H. L. (2005). Synthesis, Crystal Structure, and Nonlinear Optical Behavior of  $\beta$ -Unsubstituted meso-meso E-Vinylene-Linked Porphyrin Dimers. *Organic Letters*, 7(24), 5365-5368.
- Freindorf, M. and Gao, J. (1996). Optimization of the Lennard-Jones parameters for a combined ab initio quantum mechanical and molecular mechanical potential using the 3-21G basis set. *Journal of Computational Chemistry*, 17(4), 386-395.
- Frenette, M., Hatamimoslehabadi, M., Bellinger-Buckley, S., Laoui, S., La, J., Bag, S. . . . Yelleswarapu, C. (2014). Shining light on the dark side of imaging: excited state absorption enhancement of a bis-styryl BODIPY photoacoustic contrast agent. *Journal of the American Chemical Society*, 136(45), 15853-15856.
- Frisch, Æ., Keith, T. A. and Dennington, R. D. (2003). *GaussView reference*. Gaussian.
- Frisch, M. J., Pople, J. A. and Binkley, J. S. (1984). Self-consistent molecular orbital methods 25. Supplementary functions for Gaussian basis sets. *The Journal of Chemical Physics*, 80(7), 3265-3269.
- Fujii, H. (2002). Electronic structure and reactivity of high-valent oxo iron porphyrins. *Coordination Chemistry Reviews*, 226(1), 51-60.
- Fujimoto, K., Yorimitsu, H. and Osuka, A. (2014). Facile Preparation of  $\beta$ -Haloporphyrins as Useful Precursors of  $\beta$ -Substituted Porphyrins. *Organic Letters*, 16(3), 972-975.
- Fungo, F., Otero, L. A., Sereno, L., Silber, J. J. and Durantini, E. N. (2000). Synthesis of porphyrin dyads with potential use in solar energy conversion. *Journal of Materials Chemistry*, 10(3), 645-650.
- Galich, L., Hückstädt, H. and Homborg, H. (1998). Tetra (n-butyl) ammonium dicyanotetraphenylporphyrinatomanganate (III); crystal structure and electronic resonance Raman and absorption spectra. *Journal of Porphyrins and Phthalocyanines*, 2(1), 79-87.

- Gao, F., Bai, L., Feng, X., Tham, H. P., Zhang, R., Zhang, Y. . . . Zhao, Y. (2016). Remarkable In Vivo Nonlinear Photoacoustic Imaging Based on Near-Infrared Organic Dyes. *Small*, 12(38), 5239-5244.
- Gao, F., Bai, L., Liu, S., Zhang, R., Zhang, J., Feng, X. . . . Zhao, Y. (2017). Rationally encapsulated gold nanorods improving both linear and nonlinear photoacoustic imaging contrast in vivo. *Nanoscale*, 9(1), 79-86.
- Gao, F., Feng, X. and Zheng, Y. (2016). Advanced photoacoustic and thermoacoustic sensing and imaging beyond pulsed absorption contrast. *Journal of Optics*, 18(7), 74006-74025.
- Gaussian09, R. A. (2009). 1, MJ Frisch, GW Trucks, HB Schlegel, GE Scuseria, MA Robb, JR Cheeseman, G. Scalmani, V. Barone, B. Mennucci, GA Petersson et al., Gaussian. Inc., Wallingford CT.
- Geier III, G. R., Callinan, J. B., Dharma Rao, P. and Lindsey, J. S. (2001). A survey of acid catalysts in dipyrromethanecarbinol condensations leading to meso-substituted porphyrins. *Journal of Porphyrins and Phthalocyanines*, 5(12), 810-823.
- Geier III, G. R. and Lindsey, J. S. (2001). Investigation of porphyrin-forming reactions. Part 2. 1 Examination of the reaction course in two-step, one-flask syntheses of meso-substituted porphyrins. *Journal of the Chemical Society, Perkin Transactions*, 2(5), 687-700.
- Geier III, G. R., Littler, B. J. and Lindsey, J. S. (2001). Investigation of porphyrin-forming reactions. Part 3. 1 The origin of scrambling in dipyrromethane+ aldehyde condensations yielding trans-A 2 B 2-tetraarylporphyrins. *Journal of the Chemical Society, Perkin Transactions*, 2(5), 701-711.
- Giovannetti, R. (2012). The use of Spectrophotometry UV-Vis for the Study of Porphyrins. In Dr. Jamal Uddin (Ed.), *Macro to nano spectroscopy* (pp. 87-108). Rijeka, Croatia: InTech.
- Gnach, A., Lipinski, T., Bednarkiewicz, A., Rybka, J. and Capobianco, J. A. (2015). Upconverting nanoparticles: assessing the toxicity. *Chemical Society Reviews*, 44(6), 1561-1584.
- Goerigk, L. and Grimme, S. (2008). Calculation of electronic circular dichroism spectra with time-dependent double-hybrid density functional theory. *The Journal of Physical Chemistry A*, 113(4), 767-776.
- Gonsalves, A. M. d. A. R., Johnstone, R. A., Pereira, M. M., Shaw, J. and Sobral, A. J. d. N. (1991). Metal-assisted reactions. Part 22. Synthesis of perhalogenated porphyrins and their use as oxidation catalysts. *Tetrahedron Letters*, 32(10), 1355-1358.

- Gosper, J. J. and Ali, M. (1994). A conformationally constrained conjugated porphyrin dimer. *Journal of the Chemical Society, Chemical Communications*, (14), 1707-1708.
- Gouterman, M. (1959). Study of the effects of substitution on the absorption spectra of porphin. *The Journal of Chemical Physics*, 30(5), 1139-1161.
- Gouterman, M. (1961). Spectra of porphyrins. *Journal of Molecular Spectroscopy*, 6, 138-163.
- Gouterman, M. (1978). Optical Spectra and Electronic Structure of Porphyrins and Related Rings. In D. Dolphin (ed.), *The Porphyrins. Physical Chemistry Part 1 (Vol. 3 pp. 1-165)*. New York: Academic Press.
- Gouterman, M., Rentzepis, P. M. and Straub, K. D. (1986). Porphyrins: excited states and dynamics. *Analytical Chemistry*, 58 (14), 1438A-1438A.
- Grinstaff, M. W., Hill, M. G., Labinger, J. A. and Gray, H. B. (1994). Mechanism of catalytic oxygenation of alkanes by halogenated iron porphyrins. *Science*, 264(5163), 1311-1314.
- Gross, E. K. and Dreizler, R. M. (2013). *Density functional theory*. New York: Springer Science & Business Media.
- Guggenheim, J. A., Allen, T. J., Plumb, A., Zhang, E. Z., Rodriguez-Justo, M., Punwani, S. and Beard, P. C. (2015). Photoacoustic imaging of human lymph nodes with endogenous lipid and hemoglobin contrast. *Journal of Biomedical Optics*, 20(5), 050504-050504.
- Guilbault, G. G. (1990). *Practical fluorescence*. Second Edition, revised and expanded. New York: Marcel Dekker Inc.
- Ham, N. S. and Ruedenberg, K. (1956a). Electronic Interaction in the Free-Electron Network Model for Conjugated Systems. I. Theory. *The Journal of Chemical Physics*, 25(1), 1-13.
- Ham, N. S. and Ruedenberg, K. (1956b). Electronic Interaction in the Free-Electron Network Model for Conjugated Systems. II. Spectra of Aromatic Hydrocarbons. *The Journal of Chemical Physics*, 25(1), 13-26.
- Hambright, P. (1970). Kinetics of incorporation of Cu (II) and Zn (II) into meso-tetrapyridylporphin in the presence of Li (I). *Journal of Inorganic and Nuclear Chemistry*, 32(7), 2449-2452.
- Hambright, P. (1971). The coordination chemistry of metalloporphyrins. *Coordination Chemistry Reviews*, 6(2-3), 247-268.



- Han, Y., Fang, H., Jing, H., Sun, H., Lei, H., Lai, W. and Cao, R. (2016). Singly versus Doubly Reduced Nickel Porphyrins for Proton Reduction: Experimental and Theoretical Evidence for a Homolytic Hydrogen-Evolution Reaction. *Angewandte Chemie International Edition*, 55(18), 5457-5462.
- Harada, R., Matsuda, Y., Ōkawa, H., Miyamoto, R., Yamauchi, S. and Kojima, T. (2005). Synthesis and characterization of chromium (III) octaphenylporphyrin complexes with various axial ligands: An insight into porphyrin distortion. *Inorganica Chimica Acta*, 358(8), 2489-2500.
- Hariharan, P. C. and Pople, J. A. (1973). The influence of polarization functions on molecular orbital hydrogenation energies. *Theoretical Chemistry Accounts: Theory, Computation, and Modeling (Theoretica Chimica Acta)*, 28(3), 213-222.
- Harrach, G., Valicsek, Z. and Horváth, O. (2011). Water-soluble silver (II) and gold (III) porphyrins: The effect of structural distortion on the photophysical and photochemical behavior. *Inorganic Chemistry Communications*, 14(11), 1756-1761.
- Harriman, A. (1981). Luminescence of porphyrins and metalloporphyrins. Part 3.— Heavy-atom effects. *Journal of the Chemical Society, Faraday Transactions 2: Molecular and Chemical Physics*, 77(7), 1281-1291.
- Hartnell, R. D., Edwards, A. J. and Arnold, D. P. (2002). Peripherally-metallated porphyrins: meso- $\eta^1$ -porphyrinyl-platinum (II) complexes of 5, 15-diaryl- and 5, 10, 15-triarylporphyrins. *Journal of Porphyrins and Phthalocyanines*, 6(11), 695-707.
- Hartree, D. R. (1928). *The wave mechanics of an atom with a non-Coulomb central field. Part I. Theory and methods*. Paper presented at the Mathematical Proceedings of the Cambridge Philosophical Society.
- Hashimoto, T., Choe, Y.-K., Nakano, H. and Hirao, K. (1999). Theoretical study of the Q and B bands of free-base, magnesium, and zinc porphyrins, and their derivatives. *The Journal of Physical Chemistry A*, 103(12), 1894-1904.
- Hata, H., Shinokubo, H. and Osuka, A. (2005). Highly Regioselective Ir-Catalyzed  $\beta$ -Borylation of Porphyrins via C–H Bond Activation and Construction of  $\beta$ – $\beta$ -Linked Diporphyrin. *Journal of the American Chemical Society*, 127(23), 8264-8265.
- Hatamimoslehabadi, M., Bellinger, S., La, J., Ahmad, E., Frenette, M., Yelleswarapu, C. and Rochford, J. (2017). Correlation of Photophysical Properties with the Photoacoustic Emission for a Selection of Established Chromophores. *The Journal of Physical Chemistry C*, 121(43), 24168-24178.

- Hay, P. J. and Wadt, W. R. (1985a). Ab initio effective core potentials for molecular calculations. Potentials for K to Au including the outermost core orbitals. *The Journal of Chemical Physics*, 82(1), 299-310.
- Hay, P. J. and Wadt, W. R. (1985b). Ab initio effective core potentials for molecular calculations. Potentials for the transition metal atoms Sc to Hg. *The Journal of Chemical Physics*, 82(1), 270-283.
- Hayashi, S., Yotsukura, M., Noji, M. and Takanami, T. (2015). Bis (zinc porphyrin) as a CD-sensitive bidentate host molecule: Direct determination of absolute configuration of mono-alcohols. *Chemical Communications*, 51(55), 11068-11071.
- Head-Gordon, M., Rico, R. J., Oumi, M. and Lee, T. J. (1994). A doubles correction to electronic excited states from configuration interaction in the space of single substitutions. *Chemical Physics Letters*, 219(1-2), 21-29.
- Hehre, W. J., Ditchfield, R. and Pople, J. A. (1972). Self-consistent molecular orbital methods. XII. Further extensions of gaussian-type basis sets for use in molecular orbital studies of organic molecules. *The Journal of Chemical Physics*, 56(5), 2257-2261.
- Heijblom, M., Piras, D., Xia, W., van Hespen, J. C., Klaase, J., Van den Engh, F. . . . Manohar, S. (2012). Visualizing breast cancer using the Twente photoacoustic mammoscope: what do we learn from twelve new patient measurements? *Optics Express*, 20(11), 11582-11597.
- Hiroto, S., Miyake, Y. and Shinokubo, H. (2016). Synthesis and functionalization of porphyrins through organometallic methodologies. *Chemical Reviews*, 117(4), 2910-3043.
- Hisaki, I., Hiroto, S., Kim, K. S., Noh, S. B., Kim, D., Shinokubo, H. and Osuka, A. (2007). Synthesis of Doubly  $\beta$ -to- $\beta$  1, 3-Butadiyne-Bridged Diporphyrins: Enforced Planar Structures and Large Two-Photon Absorption Cross Sections. *Angewandte Chemie International Edition*, 46(27), 5125-5128.
- Ho, C. J. H., Balasundaram, G., Driessen, W., McLaren, R., Wong, C. L., Dinish, U. . . . Olivo, M. (2014). Multifunctional photosensitizer-based contrast agents for photoacoustic imaging. *Scientific Reports*, 4, 5342(1-6).
- Ho, I.-T., Sessler, J. L., Gambhir, S. S. and Jokerst, J. V. (2015). Parts per billion detection of uranium with a porphyrinoid-containing nanoparticle and in vivo photoacoustic imaging. *Analyst*, 140(11), 3731-3737.
- Hoard, J. (1971). Stereochemistry of hemes and other metalloporphyrins. *Science*, 174(4016), 1295-1302.

- Hoard, J. and Smith, K. (1975). *Porphyrins and metalloporphyrins*. Amsterdam: Elsevier.
- Hoelen, C., De Mul, F., Pongers, R. and Dekker, A. (1998). Three-dimensional photoacoustic imaging of blood vessels in tissue. *Optics Letters*, 23(8), 648-650.
- Hohenberg, P. and Kohn, W. (1964). Inhomogeneous electron gas. *Physical Review*, 136(3B), 864-871.
- Hohenberg, P., Kohn, W. and Sham, L. (1990). The beginnings and some thoughts on the future. *Advances in Quantum Chemistry*, 21, 7-26.
- Holten, D., Bocian, D. F. and Lindsey, J. S. (2002). Probing electronic communication in covalently linked multiporphyrin arrays. A guide to the rational design of molecular photonic devices. *Accounts of Chemical Research*, 35(1), 57-69.
- Höper, F. and Montforts, F. P. (1995). A simple synthesis of functionalized rac-chlorins via allylboration of porphyrin aldehydes. *European Journal of Organic Chemistry*, 1995(6), 1033-1038.
- Horning, T. L., Fujita, E. and Fajer, J. (1986). ESR and ENDOR of bacteriopheophytin a radicals. Implications for bacteriochlorophylls in vivo. *Journal of the American Chemical Society*, 108(2), 323-325.
- Horrocks Jr, W. D. and Hove, E. G. (1978). Water-soluble lanthanide porphyrins: shift reagents for aqueous solution. *Journal of the American Chemical Society*, 100(14), 4386-4392.
- Horváth, O., Huszánk, R., Valicsek, Z. and Lendvay, G. (2006). Photophysics and photochemistry of kinetically labile, water-soluble porphyrin complexes. *Coordination Chemistry Reviews*, 250(13), 1792-1803.
- Horváth, O., Valicsek, Z., Harrach, G., Lendvay, G. and Fodor, M. A. (2012). Spectroscopic and photochemical properties of water-soluble metalloporphyrins of distorted structure. *Coordination Chemistry Reviews*, 256(15), 1531-1545.
- Horváth, O., Valicsek, Z. and Vogler, A. (2004). Unique photoreactivity of mercury (II) 5, 10, 15, 20-tetrakis (4-sulfonatophenyl) porphyrin. *Inorganic Chemistry Communications*, 7(7), 854-857.
- Hu, S., Maslov, K. and Wang, L. V. (2011). Second-generation optical-resolution photoacoustic microscopy with improved sensitivity and speed. *Optics Letters*, 36(7), 1134-1136.

- Hudson, M. F. and Smith, K. M. (1974). Novel mercury (II) porphyrins-II: mercury (II) chelates of meso-tetraphenylporphyrin and N-methylporphyrins. *Tetrahedron Letters*, 15(26), 2223-2226.
- Huszánk, R. and Horváth, O. (2005). A heme-like, water-soluble iron (II) porphyrin: thermal and photoinduced properties, evidence for sitting-atop structure. *Chemical Communications*, (2), 224-226.
- Huynh, E., Jin, C. S., Wilson, B. C. and Zheng, G. (2014). Aggregate enhanced trimodal porphyrin shell microbubbles for ultrasound, photoacoustic, and fluorescence imaging. *Bioconjugate Chemistry*, 25(4), 796-801.
- Huynh, E., Lovell, J. F., Helfield, B. L., Jeon, M., Kim, C., Goertz, D. E. . . . Zheng, G. (2012). Porphyrin shell microbubbles with intrinsic ultrasound and photoacoustic properties. *Journal of the American Chemical Society*, 134(40), 16464-16467.
- Huzinaga, S. (1985). Basis sets for molecular calculations. *Computer Physics Reports*, 2(6), 281-339.
- Hynninen, P. H. and Lötjönen, S. (1980). Preparation of phorbin derivatives from chlorophyll mixture utilizing the principle of selective hydrolysis. *Synthesis*, 1980(07), 539-541.
- Ichiki, T., Matsuo, Y. and Nakamura, E. (2013). Photostability of a dyad of magnesium porphyrin and fullerene and its application to photocurrent conversion. *Chemical Communications*, 49(3), 279-281.
- Inhoffen, H. H., Fuhrhop, J. H., Voigt, H. and Brockmann, H. (1966). Zur weiteren Kenntnis des Chlorophylls und des Hämins, VI. Formylierung der meso-Kohlenstoffatome von Alkyl-substituierten Porphyrinen. *European Journal of Organic Chemistry*, 695(1), 133-143.
- Ishida, M., Hwang, D., Koo, Y. B., Sung, J., Kim, D. Y., Sessler, J. L. and Kim, D. (2013).  $\beta$ -(Ethynylbenzoic acid)-substituted push-pull porphyrins: DSSC dyes prepared by a direct palladium-catalyzed alkynylation reaction. *Chemical Communications*, 49(80), 9164-9166.
- Ishihara, S., Labuta, J., Van Rossom, W., Ishikawa, D., Minami, K., Hill, J. P. and Ariga, K. (2014). Porphyrin-based sensor nanoarchitectonics in diverse physical detection modes. *Physical Chemistry Chemical Physics*, 16(21), 9713-9746.
- Ivanova, Y. B., Semeikin, A., Glazunov, A. and Mamardashvili, N. Z. (2010). Pyridyl-substituted porphyrins: II. Synthesis and basic properties of dipyridylporphyrins. *Russian Journal of Organic Chemistry*, 46(6), 917-923.
- Jablonski, A. (1933). Efficiency of anti-Stokes fluorescence in dyes. *Nature*, 131(3319), 839-840.

- Jackson, A., Kenner, G. and Smith, K. (1971). Pyrroles and related compounds. Part XVII. Porphyrin synthesis through b-bilenes. *Journal of the Chemical Society C: Organic*, 502-509.
- Jacquemin, D., Perpète, E. A., Ciofini, I. and Adamo, C. (2010). Assessment of functionals for TD-DFT calculations of singlet– triplet transitions. *Journal of Chemical Theory and Computation*, 6(5), 1532-1537.
- Jacquemin, D., Perpète, E. A., Scuseria, G. E., Ciofini, I. and Adamo, C. (2008). TD-DFT performance for the visible absorption spectra of organic dyes: conventional versus long-range hybrids. *Journal of Chemical Theory and Computation*, 4(1), 123-135.
- Jaquinod, L. (2000). Functionalization of 5, 10, 15, 20-tetra-substituted porphyrins. In K. M. Kadish, K. M. Smith and R. Guillard (eds.), *The Porphyrin Handbook* (Vol. 1, pp. 45). San Diego: Academic Press.
- Jeandon, C., Bauder, C. and Callot, H. (1990). Selective acetylation of porphyrins possessing a free pyrrolic position as a tool for petroporphyrin analysis. A reinvestigation. *Energy and Fuels*, 4(6), 665-667.
- Jensen, F. (2013). Atomic orbital basis sets. *Wiley Interdisciplinary Reviews: Computational Molecular Science*, 3(3), 273-295.
- Jentzen, W., Ma, J.-G. and Shelnutt, J. A. (1998). Conservation of the conformation of the porphyrin macrocycle in hemoproteins. *Biophysical Journal*, 74(2), 753-763.
- Jentzen, W., Simpson, M., Hobbs, J., Song, X., Ema, T., Nelson, N. . . Mazzanti, M. (1995). Ruffling in a Series of Nickel (II) meso-Tetrasubstituted Porphyrins as a Model for the Conserved Ruffling of the Heme of Cytochromes c. *Journal of the American Chemical Society*, 117(45), 11085-11097.
- Jiang, J., Vairaprakash, P., Reddy, K. R., Sahin, T., Pavan, M. P., Lubian, E. and Lindsey, J. S. (2014). Hydrophilic tetracarboxy bacteriochlorins for photonics applications. *Organic & Biomolecular Chemistry*, 12(1), 86-103.
- Jiang, X.-L., Zhang, H.-L. and JIAN, W. (2006). Synthesis of diaryl ether-linked porphyrin dimers via ullmann coupling reaction. *Heterocycles*, 68(10), 2153-2160.
- Jiang, X., Gou, F. and Jing, H. (2014). Alternating copolymerization of CO<sub>2</sub> and propylene oxide catalyzed by C<sub>2</sub>v-porphyrin cobalt: Selectivity control and a kinetic study. *Journal of Catalysis*, 313, 159-167.
- Johnson, A. (1978). Synthesis of Porphyrins from 1, 19-Dideoxybiladienes-ac and 1, 19-Dideoxybilenes-b. In D.Dolphine (ed.), *The Porphyrins* (Vol. 1, pp. 235-264). New York: Academic Press.

- Johnson, A. and Kay, I. (1961). 468. The formation of porphyrins by the cyclisation of bilenes. *Journal of the Chemical Society (Resumed)*, 0, 2418-2423.
- Jokerst, J. V. (2014). Bioimaging: Sensed at the gut level. *Nature Nanotechnology*, 9(8), 569-570.
- Jradi, F. M., O'Neil, D., Kang, X., Wong, J., Szymanski, P., Parker, T. C. . . . Marder, S. R. (2015). A Step Toward Efficient Panchromatic Multi-Chromophoric Sensitizers for Dye Sensitized Solar Cells. *Chemistry of Materials*, 27(18), 6305-6313.
- Jurow, M., Schuckman, A. E., Batteas, J. D. and Drain, C. M. (2010). Porphyrins as molecular electronic components of functional devices. *Coordination Chemistry Reviews*, 254(19), 2297-2310.
- Jusélius, J. and Sundholm, D. (2000). The aromatic character of magnesium porphyrins. *The Journal of Organic Chemistry*, 65(17), 5233-5237.
- Kadish, K. M., Van Caemelbecke, E., Boulas, P., D'Souza, F., Vogel, E., Kisters, M. . . . Smith, K. M. (1993). First reversible electrogeneration of triply oxidized nickel porphyrins and porphycenes. Formation of nickel (III). pi. dications. *Inorganic Chemistry*, 32(20), 4177-4178.
- Kalyanasundaram, K. (1991). *Photochemistry of polypyridine and porphyrin complexes*. London: Academic Press.
- Kaplan, W. A., Suslick, K. S. and Scott, R. A. (1991). Core size and flexibility of metallohydroporphyrin macrocycles. Implications for F430 coordination chemistry. *Journal of the American Chemical Society*, 113(26), 9824-9827.
- Kasha, M. (1950). Characterization of electronic transitions in complex molecules. *Discussions of the Faraday Society*, 9, 14-19.
- Katritzky, A. R., He, H.-Y., Long, Q. and Wilcox, A. L. (2001). Preparation of 2, 6-dialkoxybenzaldehydes. *Arkivoc*, 3, 3-12.
- Kendrew, J., Dickerson, R., Strandberg, B., Hart, R., Davies, D., Phillips, D. and Shore, V. (1960). Structure of myoglobin: A three-dimensional Fourier synthesis at 2 Å resolution. *Nature*, 185(4711), 422-427.
- Kim, C., Cho, E. C., Chen, J., Song, K. H., Au, L., Favazza, C. . . . Xia, Y. (2010). In vivo molecular photoacoustic tomography of melanomas targeted by bioconjugated gold nanocages. *ACS Nano*, 4(8), 4559-4564.
- Kinoshita, H., Tanaka, S., Nishimori, N., Dejima, H. and Inomata, K. (1992). Synthesis of 2-(substituted methyl)-3, 4-disubstituted pyrroles and their conversion into the

- corresponding porphyrins. *Bulletin of the Chemical Society of Japan*, 65(10), 2660-2667.
- Knopoff, L. and Shapiro, J. (1970). Pseudo-Grüneisen parameter for liquids. *Physical Review B*, 1(10), 3893-3895.
- Kohn, W. and Sham, L. (1965). Self-consistent equations including exchange and correlation effects. *Physical Review*, 140, A1133-A1138.
- Kowalski, K., Dean, D., Hjorth-Jensen, M., Papenbrock, T. and Piecuch, P. (2004). Coupled cluster calculations of ground and excited states of nuclei. *Physical Review Letters*, 92(13), 132501(1-4).
- Kruglik, S. G., Apanasevich, P. A., Chirvony, V. S., Kvach, V. V. and Orlovich, V. A. (1995). Resonance Raman, CARS, and picosecond absorption spectroscopy of copper porphyrins: The evidence for the exciplex formation with oxygen-containing solvent molecules. *The Journal of Physical Chemistry*, 99(10), 2978-2995.
- Kumar, R. and Sankar, M. (2014). Synthesis, Spectral, and Electrochemical Studies of Electronically Tunable  $\beta$ -Substituted Porphyrins with Mixed Substituent Pattern. *Inorganic Chemistry*, 53(24), 12706-12719.
- Kuster, W. (1912). Information on bilirubine and haemine. *Hoppe-Seylers Zeitschrift Fur Physiologische Chemie*, 82(6), 463-483.
- Ladomenou, K., Nikolaou, V., Charalambidis, G. and Coutsolelos, A. G. (2016). “Click”-reaction: An alternative tool for new architectures of porphyrin based derivatives. *Coordination Chemistry Reviews*, 306, 1-42.
- Lakowicz, J. (1999a). *Principles of fluorescence spectroscopy*. New York: Kluwer Academic Plenum.
- Lakowicz, J. R. (1999b). Introduction to fluorescence. *Principles of Fluorescence Spectroscopy* (ed. 3<sup>rd</sup>, pp. 1-23). New York: Springer.
- Lash, T. D. (1996). Porphyrin synthesis by the “3+ 1” approach: new applications for an old methodology. *Chemistry-A European Journal*, 2(10), 1197-1200.
- Lash, T. D. (2000). Carbaporphyrinoids: Taking the Heterocycle Out of Nature's [18] Annulene. *Synlett*, 2000(03), 279-295.
- Le Maux, P., Bahri, H. and Simonneaux, G. (1991). Molecular recognition of racemic phosphines by a chiral ruthenium porphyrin. *Journal of the Chemical Society, Chemical Communications*, (19), 1350-1352.

- Le Maux, P., Bahri, H. and Simonneaux, G. (1993). Synthesis and stereochemical studies of chiral ruthenium porphyrins. *Tetrahedron*, 49(7), 1401-1408.
- Leach, A. R. (2001). *Molecular modelling: principles and applications*. (ed. 2<sup>nd</sup>). United Kingdom: Prentice Hall Harlow.
- Lee, C., Yang, W. and Parr, R. (1988a). Density-functional exchange-energy approximation with correct asymptotic behaviour. *Physical Review B*, 37, 785-789.
- Lee, C., Yang, W. and Parr, R. G. (1988b). Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Physical Review B*, 37(2), 785-789.
- Lewars, E. G. (2016). *Computational chemistry: introduction to the theory and applications of molecular and quantum mechanics*. Boston: Springer.
- Lewinski, N., Colvin, V. and Drezek, R. (2008). Cytotoxicity of nanoparticles. *Small*, 4(1), 26-49.
- Liao, M.-S., Watts, J. D. and Huang, M.-J. (2005). Effects of peripheral substituents and axial ligands on the electronic structure and properties of cobalt porphyrins. *The Journal of Physical Chemistry A*, 109(51), 11996-12005.
- Liao, M.-S., Watts, J. D. and Huang, M.-J. (2006). DFT/TDDFT study of lanthanide(III) mono- and bisporphyrin complexes. *The Journal of Physical Chemistry A*, 110(48), 13089-13098.
- Licha, K., Riefke, B., Ntziachristos, V., Becker, A., Chance, B. and Semmler, W. (2000). Hydrophilic Cyanine Dyes as Contrast Agents for Near-infrared Tumor Imaging: Synthesis, Photophysical Properties and Spectroscopic In vivo Characterization ¶. *Photochemistry and Photobiology*, 72(3), 392-398.
- Lin, J. J., Gerzevske, K. R., Liddell, P. A., Senge, M. O., Olmstead, M. M., Khoury, R. G. . . . Smith, K. M. (1997). Metal-catalyzed oxidative cyclizations of a, c-biladiene salts bearing 1-and/or 19-arylmethyl substituents: Macrocyclic products and their chemistry. *The Journal of Organic Chemistry*, 62(13), 4266-4276.
- Lin, V. S.-Y., DiMagno, S. G. and Therien, M. J. (1994). Highly conjugated, acetylenyl bridged porphyrins: new models for light-harvesting antenna systems. *Science*, 264(5162), 1105-1112.
- Lindsey, J. S., Schreiman, I. C., Hsu, H. C., Kearney, P. C. and Marguerettaz, A. M. (1987). Rothmund and Adler-Longo reactions revisited: synthesis of tetraphenylporphyrins under equilibrium conditions. *The Journal of Organic Chemistry*, 52(5), 827-836.



- Lindsey, J. S. and Woodford, J. N. (1995). A simple method for preparing magnesium porphyrins. *Inorganic Chemistry*, 34(5), 1063-1069.
- Littler, B. J., Miller, M. A., Hung, C.-H., Wagner, R. W., O'Shea, D. F., Boyle, P. D. and Lindsey, J. S. (1999). Refined synthesis of 5-substituted dipyrromethanes. *The Journal of Organic Chemistry*, 64(4), 1391-1396.
- Liu, C., Shen, D.-M. and Chen, Q.-Y. (2007). Practical and Efficient Synthesis of Various meso-Functionalized Porphyrins via Simple Ligand-Free Nickel-Catalyzed C–O, C–N, and C–C Cross-Coupling Reactions. *The Journal of Organic Chemistry*, 72(8), 2732-2736.
- Liu, P. P., Feng, Y. Q., Gu, C. Z., Meng, S. X. and Zhang, B. (2012). The facile synthesis of 5-formylporphyrin. *Chinese Chemical Letters*, 23(5), 505-508.
- Liu, X., Yang, G., Zhang, L., Liu, Z., Cheng, Z. and Zhu, X. (2016). Photosensitizer cross-linked nano-micelle platform for multimodal imaging guided synergistic photothermal/photodynamic therapy. *Nanoscale*, 8(33), 15323-15339.
- Ljungdahl, T., Pettersson, K., Albinsson, B. and Mårtensson, J. (2006). Solvent and base dependence of copper-free palladium-catalyzed cross-couplings between terminal alkynes and aryl iodides: Development of efficient conditions for the construction of gold (III)/free-base porphyrin dimers. *The Journal of Organic Chemistry*, 71(4), 1677-1687.
- Locos, O. B. and Arnold, D. P. (2006). The Heck reaction for porphyrin functionalisation: synthesis of meso-alkenyl monoporphyrins and palladium-catalysed formation of unprecedented meso- $\beta$  ethene-linked diporphyrins. *Organic & Biomolecular Chemistry*, 4(5), 902-916.
- Longo, F. R. (1979). *Porphyrin chemistry advances*. New York: Ann Arbor Science Publishers Inc.
- Lovell, J. F., Jin, C. S., Huynh, E., Jin, H., Kim, C., Rubinstein, J. L. . . . Zheng, G. (2011). Porphysome nanovesicles generated by porphyrin bilayers for use as multimodal biophotonic contrast agents. *Nature Materials*, 10(4), 324-332.
- Luciano, M., Erfanzadeh, M., Zhou, F., Zhu, H., Bornhütter, T., Röder, B. . . . Brückner, C. (2016). In vivo photoacoustic tumor tomography using a quinoline-annulated porphyrin as NIR molecular contrast agent. *Organic & Biomolecular Chemistry*, 15(4), 972-983.
- Luguya, R., Jaquinod, L., Fronczek, F. R., Vicente, M. G. H. and Smith, K. M. (2004). Synthesis and reactions of meso-(p-nitrophenyl) porphyrins. *Tetrahedron*, 60(12), 2757-2763.

- Luo, S., Zhang, E., Su, Y., Cheng, T. and Shi, C. (2011). A review of NIR dyes in cancer targeting and imaging. *Biomaterials*, 32(29), 7127-7138.
- Luthra, A., Denisov, I. G. and Sligar, S. G. (2011). Spectroscopic features of cytochrome P450 reaction intermediates. *Archives of Biochemistry and Biophysics*, 507(1), 26-35.
- Lysenko, A. B., Thamyongkit, P., Schmidt, I., Diers, J. R., Bocian, D. F. and Lindsey, J. S. (2006). Diverse porphyrin dimers as candidates for high-density charge-storage molecules. *Journal of Porphyrins and Phthalocyanines*, 10(01), 22-32.
- MacDonald, F., Arsenault, P. and Bullock, E. (1960). Pyrromethanes and porphyrins therefrom. *Journal of American Chemical Society*, 82, 4384-4389.
- Maji, S. K., Sreejith, S., Joseph, J., Lin, M., He, T., Tong, Y. . . . Zhao, Y. (2014). Upconversion nanoparticles as a contrast agent for photoacoustic imaging in live mice. *Advanced Materials*, 26(32), 5633-5638.
- Mallidi, S., Larson, T., Tam, J., Joshi, P. P., Karpouk, A., Sokolov, K. and Emelianov, S. (2009). Multiwavelength photoacoustic imaging and plasmon resonance coupling of gold nanoparticles for selective detection of cancer. *Nano Letters*, 9(8), 2825-2831.
- Mamardashvili, N. Z. and Golubchikov, O. A. (2000). The synthesis of porphyrins from dipyrrolylmethanes. *Russian Chemical Reviews*, 69(4), 307-323.
- Markovac, A. and MacDonald, S. (1965). Syntheses with 5-dibromomethyl- and 5-formyl-pyrromethenes. *Canadian Journal of Chemistry*, 43(12), 3364-3371.
- Marques, M. A. and Gross, E. K. (2003). Time-dependent density functional theory. *Lecture Notes In Physics-New York Then Berlin*, 144-184.
- Marques, M. A., Maitra, N. T., Nogueira, F. M., Gross, E. K. and Rubio, A. (2012). *Fundamentals of time-dependent density functional theory*. New York: Springer Science & Business Media.
- Márquez, N., Ysambertt, F. and De La Cruz, C. (1999). Three analytical methods to isolate and characterize vanadium and nickel porphyrins from heavy crude oil. *Analytica Chimica Acta*, 395(3), 343-349.
- Marten, B., Kim, K., Cortis, C., Friesner, R. A., Murphy, R. B., Ringnalda, M. N. . . . Honig, B. (1996). New model for calculation of solvation free energies: correction of self-consistent reaction field continuum dielectric theory for short-range hydrogen-bonding effects. *The Journal of Physical Chemistry*, 100(28), 11775-11788.

- Matano, Y., Fujii, D., Shibano, T., Furukawa, K., Higashino, T., Nakano, H. and Imahori, H. (2014). Covalently Linked 5, 15-Diazaporphyrin Dimers: Promising Scaffolds for a Highly Conjugated Azaporphyrin  $\pi$  System. *Chemistry-A European Journal*, 20(12), 3342-3349.
- Mathew, S., Yella, A., Gao, P., Humphry-Baker, R., Curchod, B. F., Ashari-Astani, N. . . . Grätzel, M. (2014). Dye-sensitized solar cells with 13% efficiency achieved through the molecular engineering of porphyrin sensitizers. *Nature Chemistry*, 6(3), 242-247.
- McGlynn, S., Azumi, T. and Kasha, M. (1964). External Heavy-Atom Spin—Orbital Coupling Effect. V. Absorption Studies of Triplet States. *The Journal of Chemical Physics*, 40(2), 507-515.
- Milgrom, L. R. and Warren, M. J. (1997). *The colours of life: an introduction to the chemistry of porphyrins and related compounds*. Oxford: Oxford University Press.
- Milroy, J. A. (1918). Observations on some Metallic Compounds of Haematoporphyrin. *Biochemical Journal*, 12(4), 318-338.
- Minaev, B. and Lindgren, M. (2009). Vibration and fluorescence spectra of porphyrin-cored bis (methylol)-propionic acid dendrimers. *Sensors*, 9(3), 1937-1966.
- Mitra, K., Singha, A. and Dey, A. (2016). Iron porphyrins with a hydrogen bonding cavity: Effect of weak interactions on their electronic structure and reactivity. *Dalton Transactions*, 45(47), 18796-18802.
- Mohr, B., Enkelmann, V. and Wegner, G. (1994). Synthesis of alkyl- and alkoxy-substituted benziles and oxidative coupling to tetraalkoxyphenanthrene-9, 10-diones. *Journal of Organic Chemistry*, 59(3), 635-638.
- Monteiro, C. J., Pereira, M. M., Vicente, M. G. H. and Arnaut, L. G. (2012). Photophysical properties of unsymmetric meso-substituted porphyrins synthesized via the Suzuki coupling reaction. *Tetrahedron*, 68(42), 8783-8788.
- Montgomery Jr, J. A., Frisch, M. J., Ochterski, J. W. and Petersson, G. A. (1999). A complete basis set model chemistry. VI. Use of density functional geometries and frequencies. *The Journal of Chemical Physics*, 110(6), 2822-2827.
- Mori, G., Shinokubo, H. and Osuka, A. (2008). Highly selective Ir-catalyzed direct sixfold borylation of peripheral aromatic substituents on hexakisaryl-substituted [28] hexaphyrin (1.1. 1.1. 1.1). *Tetrahedron Letters*, 49(13), 2170-2172.
- Naik, A., Rubbiani, R., Gasser, G. and Spingler, B. (2014). Visible-Light-Induced Annihilation of Tumor Cells with Platinum–Porphyrin Conjugates. *Angewandte Chemie International Edition*, 53(27), 6938-6941.

- Nakamura, M., Ikezaki, A. and Takahashi, M. (2013). Metal-porphyrin orbital interactions in paramagnetic iron complexes having planar and deformed porphyrin ring. *Journal of the Chinese Chemical Society*, 60(1), 9-21.
- Nappa, M. and Valentine, J. S. (1978). The influence of axial ligands on metalloporphyrin visible absorption spectra. Complexes of tetraphenylporphinatozinc. *Journal of the American Chemical Society*, 100(16), 5075-5080.
- Naruta, Y., Tani, F. and Maruyama, K. (1992). Meso-perfluorination of porphyrins with N-fluoropyridinium triflate. *Tetrahedron Letters*, 33(8), 1069-1072.
- Natali, M., Luisa, A., Iengo, E. and Scandola, F. (2014). Efficient photocatalytic hydrogen generation from water by a cationic cobalt (II) porphyrin. *Chemical Communications*, 50(15), 1842-1844.
- Nazeeruddin, M. K., De Angelis, F., Fantacci, S., Selloni, A., Viscardi, G., Liska, P. . . . Grätzel, M. (2005). Combined experimental and DFT-TDDFT computational study of photoelectrochemical cell ruthenium sensitizers. *Journal of the American Chemical Society*, 127(48), 16835-16847.
- Nelson, J. S., Roberts, W. G. and Berns, M. W. (1987). In vivo studies on the utilization of mono-L-aspartyl chlorin (NPe6) for photodynamic therapy. *Cancer Research*, 47(17), 4681-4685.
- Nemykin, V. N. and Hadt, R. G. (2010). Interpretation of the UV– vis Spectra of the meso (Ferrocenyl)-Containing Porphyrins using a TDDFT Approach: Is Gouterman's Classic Four-Orbital Model Still in Play? *The Journal of Physical Chemistry A*, 114(45), 12062-12066.
- Neugebauer, J., Baerends, E. J. and Nooijen, M. (2005). Vibronic structure of the permanganate absorption spectrum from time-dependent density functional calculations. *The Journal of Physical Chemistry A*, 109(6), 1168-1179.
- Neya, S., Quan, J., Hoshino, T., Hata, M. and Funasaki, N. (2004). Convenient synthesis of porphine from  $\beta$ -tetra (tert-butyl) porphyrin. *Tetrahedron Letters*, 45(47), 8629-8630.
- Neya, S., Yodo, H. and Funasaki, N. (1993). Convenient synthesis of porphine. *Journal of Heterocyclic Chemistry*, 30(2), 549-550.
- Nielsen, A. B. and Holder, A. J. (1998). *GaussView. User's reference*. Gaussian, Incorporated.
- Ol'shevskaya, V. A., Zaytsev, A. V., Savchenko, A. N., Shtil, A. A., Cheong, C.-S. and Kalinin, V. N. (2007). Boronated porphyrins and chlorins as potential anticancer drugs. *Bulletin of the Korean Chemical Society*, 28(11), 1910-1916.

- Ol'shevskaya, V. A., Zaitsev, A. V., Luzgina, V. N., Kondratieva, T. T., Ivanov, O. G., Kononova, E. G. . . . Hofmann, J. (2006). Novel boronated derivatives of 5, 10, 15, 20-tetraphenylporphyrin: synthesis and toxicity for drug-resistant tumor cells. *Bioorganic and Medicinal Chemistry*, 14(1), 109-120.
- Ono, N., Kawamura, H., Bougauchi, M. and Maruyama, K. (1990). Porphyrin synthesis from nitrocompounds. *Tetrahedron*, 46(21), 7483-7496.
- Ostrowski, S. (2005). Preparation of 2-Nitro-5, 10, 15, 20-tetraphenylporphyrin Zinc (II) Complex-post scriptum to Electrophilic Nitration of Porphyrins at the beta-Pyrrolic Position. *Polish Journal of Chemistry*, 79(7), 1169-1172.
- Osuka, A. and Shimidzu, H. (1997). meso, meso-Linked Porphyrin Arrays. *Angewandte Chemie International Edition*, 36(1-2), 135-137.
- Paine III, J. B., Kirshner, W. B., Moskowitz, D. W. and Dolphin, D. (1976). An improved synthesis of octaethylporphyrin. *The Journal of Organic Chemistry*, 41(24), 3857-3860.
- Paproski, R. J., Forbrich, A., Huynh, E., Chen, J., Lewis, J. D., Zheng, G. and Zemp, R. J. (2016). Porphyrin Nanodroplets: Sub-micrometer Ultrasound and Photoacoustic Contrast Imaging Agents. *Small*, 12(3), 371-380.
- Pereira, Simões, Tomé and Almeida Paz. (2016). Porphyrin-Based Metal-Organic Frameworks as Heterogeneous Catalysts in Oxidation Reactions. *Molecules (Basel, Switzerland)*, 21(10), 1348(1-19).
- Pereira, A. M., Alonso, C. M., Neves, M. G., Tomé, A. C., Silva, A. M., Paz, F. A. and Cavaleiro, J. A. (2008). A New Synthetic Approach to N-Arylquinolino [2, 3, 4-at] porphyrins from  $\beta$ -Arylaminoporphyrins. *The Journal of Organic Chemistry*, 73(18), 7353-7356.
- Pereira, M. M., Muller, G., Ordinas, J. I., Azenha, M. E. and Arnaut, L. G. (2002). Synthesis of vinylated 5, 10, 15, 20-tetraphenylporphyrins via Heck-type coupling reaction and their photophysical properties. *Journal of the Chemical Society, Perkin Transactions*, 2(9), 1583-1588.
- Perpète, E. A., Wathélet, V., Preat, J., Lambert, C. and Jacquemin, D. (2006). Toward a theoretical quantitative estimation of the  $\lambda_{\text{max}}$  of anthraquinones-based dyes. *Journal of Chemical Theory and Computation*, 2(2), 434-440.
- Perutz, M., Rossmann, M., Cullis, A., Muirhead, H., Will, G. and North, A. (1960). *Structure of hemoglobin*. Paper presented at the Brookhaven Symp Biol.
- Pietro, W. J., Francel, M. M., Hehre, W. J., DeFrees, D. J., Pople, J. A. and Binkley, J. S. (1982). Self-consistent molecular orbital methods. 24. Supplemented small split-

- valence basis sets for second-row elements. *Journal of the American Chemical Society*, 104(19), 5039-5048.
- Pietro, W. J. and Hehre, W. J. (1983). Molecular orbital theory of the properties of inorganic and organometallic compounds. 3. STO-3G basis sets for first-and second-row transition metals. *Journal of Computational Chemistry*, 4(2), 241-251.
- Poland, C. A., Duffin, R., Kinloch, I., Maynard, A., Wallace, W. A. H., Seaton, A. . . . Donaldson, K. (2008). Carbon nanotubes introduced into the abdominal cavity of mice show asbestos-like pathogenicity in a pilot study. *Nature Nanotechnology*, 3(7), 423-428.
- Poncet, J.-L., Barbe, J.-M., Guillard, R., Oumous, H., Lecomte, C. and Protas, J. (1982). Vanadium (II) porphyrin complexes: synthesis and characterization. Crystal structure of 2,3,7,8,12,13,17,18-octaethylporphinatobis-(dimethylphenylphosphine)vanadium(II). *Journal of the Chemical Society, Chemical Communications*, (24), 1421-1422.
- Prasath, R., Butcher, R. and Bhavana, P. (2012). Nitrothienylporphyrins: Synthesis, crystal structure and, the effect of position and number of nitro groups on the spectral and electrochemical properties. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 87, 258-264.
- Prins, L. J., Reinhoudt, D. N. and Timmerman, P. (2001). Noncovalent synthesis using hydrogen bonding. *Angewandte Chemie International Edition*, 40(13), 2382-2426.
- Prushan, M. (2005). Absorption and Fluorescence Spectroscopy of Tetraphenylporphyrin and Metallo-Tetraphenylporphyrin (pp. 1-9).
- Radon, M. (2015). Role of Spin States in Nitric Oxide Binding to Cobalt (II) and Manganese (II) Porphyrins. Is Tighter Binding Always Stronger? *Inorganic Chemistry*, 54(12), 5634-5645.
- Rao, P. D., Dhanalekshmi, S., Littler, B. J. and Lindsey, J. S. (2000). Rational syntheses of porphyrins bearing up to four different meso substituents. *The Journal of Organic Chemistry*, 65(22), 7323-7344.
- Rassolov, V. A., Pople, J. A., Ratner, M. A. and Windus, T. L. (1998). 6-31G\* basis set for atoms K through Zn. *The Journal of Chemical Physics*, 109(4), 1223-1229.
- Ricciardi, G., Rosa, A., Baerends, E. J. and van Gisbergen, S. A. (2002). Electronic structure, chemical bond, and optical spectra of metal bis (porphyrin) complexes: a DFT/TDDFT study of the bis (porphyrin) M (IV)(M= Zr, Ce, Th) series. *Journal of the American Chemical Society*, 124(41), 12319-12334.

- Rieffel, J., Chen, F., Kim, J., Chen, G., Shao, W., Shao, S. . . . Nickles, R. J. (2015). Hexamodal Imaging with Porphyrin-Phospholipid-Coated Upconversion Nanoparticles. *Advanced Materials*, 27(10), 1785-1790.
- Rillema, D., Nagle, J., Barringer Jr, L. and Meyer, T. (1981). Redox properties of metalloporphyrin excited states, lifetimes, and related properties of a series of para-substituted tetraphenylporphine carbonyl complexes of ruthenium (II). *Journal of the American Chemical Society*, 103(1), 56-62.
- Risch, N., Gauler, R. and Keuper, R. (1999). Synthesis of porphyrin dimers using a Heck-type coupling reaction with bisacrylates. *Tetrahedron Letters*, 40(15), 2925-2926.
- Rothemund, P. (1935). Formation of porphyrins from pyrrole and aldehydes. *Journal of the American Chemical Society*, 57(10), 2010-2011.
- Rothemund, P. (1939). Porphyrin studies. III. 1 The structure of the Porphine<sup>2</sup> ring system. *Journal of the American Chemical Society*, 61(10), 2912-2915.
- Rothemund, P. and Menotti, A. R. (1941). Porphyrin Studies. IV. 1 The Synthesis of  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ -Tetraphenylporphine. *Journal of the American Chemical Society*, 63(1), 267-270.
- Rousseau, K. and Dolphin, D. (1974). A purification of meso-tetraphenylporphyrin. *Tetrahedron Letters*, 15(48), 4251-4254.
- Rubio, M., Roos, B. O., Serrano-Andrés, L. and Merchán, M. (1999). Theoretical study of the electronic spectrum of magnesium-porphyrin. *The Journal of Chemical Physics*, 110(15), 7202-7209.
- Runge, E. and Gross, E. K. (1984). Density-functional theory for time-dependent systems. *Physical Review Letters*, 52(12), 997-1000.
- Runge, S. and Senge, M. O. (1999). Reaction of  $\beta$ -formylporphyrins with organometallic reagents—A facile method for the preparation of porphyrins with exocyclic double bonds. *Tetrahedron*, 55(34), 10375-10390.
- Rybicka-Jasińska, K., Shan, W., Zawada, K., Kadish, K. M. and Gryko, D. (2016). Porphyrins as photoredox catalysts: Experimental and theoretical studies. *Journal of the American Chemical Society*, 138(47), 15451-15458.
- Saltsman, I., Goldberg, I., Balasz, Y. and Gross, Z. (2007). Porphine and pyrrole-substituted porphyrin from cyclocondensation of tripyrrane with mono-substituted pyrroles. *Tetrahedron Letters*, 48(2), 239-244.
- Scales, L. (1985). *Introduction to non-linear optimization*. New York: Springer-Verlag New York, Inc.

- Schaberle, F. A., Arnaut, L. G., Serpa, C., Silva, E. F., Pereira, M. M., Abreu, A. R. and Simões, S. (2010). Infrared absorbing dyes tailored for detection and therapy of solid tumors. *Laser Applications in Life Sciences*, ed. M. Kinnunen and R. Myllylä, Oulu, 73760X.
- Schaberle, F. A., Reis, L. A., Sá, G. F., Serpa, C., Abreu, A. R., Pereira, M. M. and Arnaut, L. G. (2011). *Multi-spectral photoacoustic mapping of bacteriochlorins diffusing through the skin: exploring a new PAT contrast agent*. Paper presented at the European Conference on Biomedical Optics.
- Scheidt, W. R., Duval, H. F., Neal, T. J. and Ellison, M. K. (2000). Intrinsic structural distortions in five-coordinate (nitrosyl) iron (II) porphyrinate derivatives. *Journal of the American Chemical Society*, 122(19), 4651-4659.
- Schlegel, H. B. (1989). Some practical suggestions for optimizing geometries and locating transition states. *New Theoretical Concepts for Understanding Organic Reactions*, 267, 33-53.
- Schrödinger, E. (1926). An undulatory theory of the mechanics of atoms and molecules. *Physical Review*, 28(6), 1049-1070.
- Schwarz, M., Buehler, A., Aguirre, J. and Ntziachristos, V. (2016). Three-dimensional multispectral optoacoustic mesoscopy reveals melanin and blood oxygenation in human skin in vivo. *Journal of Biophotonics*, 9(1-2), 55-60.
- Scolaro, L. M., Romeo, A., Castriciano, M. A. and Micali, N. (2005). Unusual optical properties of porphyrin fractal J-aggregates. *Chemical Communications*, (24), 3018-3020.
- Screen, T. E., Blake, I. M., Rees, L. H., Clegg, W., Borwick, S. J. and Anderson, H. L. (2002). Making conjugated connections to porphyrins: a comparison of alkyne, alkene, imine and azo links. *Journal of the Chemical Society, Perkin Transactions*, 1(3), 320-329.
- Şen, P., Hirel, C., Andraud, C., Aronica, C., Bretonnière, Y., Mohammed, A. . . . Baryshnikov, G. (2010). Fluorescence and FTIR spectra analysis of trans-A2B2-substituted di-and tetra-phenyl porphyrins. *Materials*, 3(8), 4446-4475.
- Senge, M. O., Fazekas, M., Notaras, E. G., Blau, W. J., Zawadzka, M., Locos, O. B. and Ni Mhuirheartaigh, E. M. (2007). Nonlinear optical properties of porphyrins. *Advanced Materials*, 19(19), 2737-2774.
- Sergeeva, N. N., Scala, A., Bakar, M. A., O’Riordan, G., O’Brien, J., Grassi, G. and Senge, M. O. (2009). Synthesis of stannyl porphyrins and porphyrin dimers via stille coupling and their <sup>119</sup>Sn NMR and fluorescence properties. *The Journal of Organic Chemistry*, 74(18), 7140-7147.



- Serpa, C., Schabauer, J., Piedade, A. P., Monteiro, C. J., Pereira, M. M., Douglas, P. . . . Arnaut, L. G. (2008). Photoacoustic measurement of electron injection efficiencies and energies from excited sensitizer dyes into nanocrystalline TiO<sub>2</sub> films. *Journal of the American Chemical Society*, 130(28), 8876-8877.
- Sessler, J. L., Johnson, M. R. and Lynch, V. (1987). Synthesis and crystal structure of a novel tripyrrane-containing porphyrinogen-like macrocycle. *The Journal of Organic Chemistry*, 52(19), 4394-4397.
- Sessler, J. L., Mozaffari, A. and Johnson, M. R. (1991). 3, 4-Diethylpyrrole and 2, 3, 7, 8, 12, 13, 17, 18-Octaethylporphyrin. *Organic Syntheses*, 68-68.
- Sessler, J. L. and Tomat, E. (2007). Transition-metal complexes of expanded porphyrins. *Accounts of Chemical Research*, 40(5), 371-379.
- Seybold, P. G. and Gouterman, M. (1969). Porphyrins: XIII: Fluorescence spectra and quantum yields. *Journal of Molecular Spectroscopy*, 31(1-13), 1-13.
- Shanmugathan, S., Johnson, C. K., Edwards, C., Matthews, E. K., Dolphin, D. and Boyle, R. W. (2000). Regioselective halogenation and palladium-catalysed couplings on 5, 15-diphenylporphyrin. *Journal of Porphyrins and Phthalocyanines*, 4(3), 228-232.
- Sharma, R. K., Ahuja, G. and Sidhwani, I. T. (2009). A new one pot and solvent-free synthesis of nickel porphyrin complex. *Green Chemistry Letters and Reviews*, 2(2), 101-105.
- Shea, K. M., Jaquinod, L. and Smith, K. M. (1998). Dihydroporphyrin synthesis: new methodology. *The Journal of Organic Chemistry*, 63(20), 7013-7021.
- Shelby, M. L., Mara, M. W. and Chen, L. X. (2014). New insight into metalloporphyrin excited state structures and axial ligand binding from X-ray transient absorption spectroscopic studies. *Coordination Chemistry Reviews*, 277, 291-299.
- Sheldon, R. A. (1994). *Metalloporphyrins in catalytic oxidations*. New York: CRC Press.
- Shelnutt, J. and Medforth, C. (1998). Nonplanar porphyrins and their significance in proteins. *Chemical Society Reviews*, 27(1), 31-42.
- Shelnutt, J. and Ortiz, V. (1985). Substituent effects on the electronic structure of metalloporphyrins: a quantitative analysis in terms of four-orbital-model parameters. *The Journal of Physical Chemistry*, 89(22), 4733-4739.
- Shelton, R. L. and Applegate, B. E. (2010). Ultrahigh resolution photoacoustic microscopy via transient absorption. *Biomedical Optics Express*, 1(2), 676-686.

- Shen, D. M., Liu, C. and Chen, Q. Y. (2007). Synthesis and Versatile Reactions of  $\beta$ -Azidotetraarylporphyrins. *European Journal of Organic Chemistry*, 2007(9), 1419-1422.
- Shi, B. and Boyle, R. W. (2002). Synthesis of unsymmetrically substituted meso-phenylporphyrins by Suzuki cross coupling reactions. *Journal of the Chemical Society, Perkin Transactions, 1*(11), 1397-1400.
- ShingáChan, K. and Thomas, C. (1994). Synthesis of  $\beta$ -aryl substituted porphyrins by palladium-catalysed cross-coupling reactions. *Journal of the Chemical Society, Chemical Communications*, (3), 271-272.
- Shoji, Y., Tashiro, K. and Aida, T. (2004). Selective extraction of higher fullerenes using cyclic dimers of zinc porphyrins. *Journal of the American Chemical Society*, 126(21), 6570-6571.
- Siri, O., Jaquinod, L. and Smith, K. M. (2000). Coplanar conjugated  $\beta$ -nitroporphyrins and some aspects of nitration of porphyrins with  $\text{N}_2\text{O}_4$ . *Tetrahedron Letters*, 41(19), 3583-3587.
- Slater, J. C. (1930). Atomic shielding constants. *Physical Review*, 36(1), 57-64.
- Smith, K., Lindsey, J., Vicente, M., Jaquinod, L., Senge, M., Callot, H. and Ocampo, R. (2000). Cyclizations of a,c-Biladiene Salts to Give Porphyrins and Their Derivatives. In K. M. Kadish, K. M. Smith and R. Guillard (eds.), *The Porphyrin Handbook—Synthesis and Organic Chemistry* (Vol. 1, pp. 119-148). San Diego: Academic Press.
- Smith, K. M. (2000). Strategies for the synthesis of octaalkylporphyrin systems. In K. M. Kadish, K. M. Smith and R. Guillard (eds.), *The porphyrin handbook—Synthesis and Organic Chemistry* (Vol. 1, pp. 119-148). San Diego: Academic Press.
- Smith K. M. (Ed.). (1975). *Porphyrins and Metalloporphyrins*, Amsterdam, Elsevier
- Smith, K. M., Bisset, G. M. and Tabba, H. D. (1982). Polyformylation of copper (II) complexes of octa-alkylporphyrins. *Journal of the Chemical Society, Perkin Transactions, 1*, 581-585.
- Smith, K. M., Goff, D. A. and Abraham, R. J. (1984). The NMR spectra of porphyrins. 27—proton NMR spectra of chlorophyll-a and pheophytin-a. *Magnetic Resonance in Chemistry*, 22(12), 779-783.
- Smith, N. W. and Dzyuba, S. V. (2010). Efficient nitration of meso-tetraphenylporphyrin with nitronium tetrafluoroborate. *Arkivoc*, 7, 10-18.

- Sommer, J. R., Shelton, A. H., Parthasarathy, A., Ghiviriga, I., Reynolds, J. R. and Schanze, K. S. (2011). Photophysical properties of near-infrared phosphorescent  $\pi$ -extended platinum porphyrins. *Chemistry of Materials*, 23(24), 5296-5304.
- Sondheimer, F., Wolovsky, R. and Amiel, Y. (1962). Unsaturated Macrocyclic Compounds. XXIII. 1 The Synthesis of the Fully Conjugated Macrocyclic Polyenes Cyclooctadecanonaene ([18] Annulene), 2 Cyclotetracosadodecaene ([24] Annulene), and Cyclotriaccontapentadecaene ([30] Annulene). *Journal of the American Chemical Society*, 84(2), 274-284.
- Soret, J. (1883). Analyse spectrale: Sur le spectre d'absorption du sang dans la partie violette et ultra-violette. *Comptes Rendus Chimie*, 97, 1269-1273.
- Sreejith, S., Huong, T. T. M., Borah, P. and Zhao, Y. (2015). Organic-inorganic nanohybrids for fluorescence, photoacoustic and Raman bioimaging. *Science Bulletin*, 60(7), 665-678.
- Sreejith, S., Joseph, J., Lin, M., Menon, N. V., Borah, P., Ng, H. J. . . . Zhao, Y. (2015). Near-infrared squaraine dye encapsulated micelles for in vivo fluorescence and photoacoustic bimodal imaging. *ACS Nano*, 9(6), 5695-5704.
- Srivastava, V., Gusain, D. and Sharma, Y. C. (2015). Critical review on the toxicity of some widely used engineered nanoparticles. *Industrial & Engineering Chemistry Research*, 54(24), 6209-6233.
- Stepien, M. and Latos-Grazynski, L. (2008). Aromaticity and tautomerism in porphyrins and porphyrinoids. In T. M. Krygowski and M. K. Cyranski (eds.), *Aromaticity in Heterocyclic Compounds* (Vol.19, pp. 84-146). Berlin: Springer.
- Strohm, E. M., Berndl, E. S. and Kolios, M. C. (2013). High frequency label-free photoacoustic microscopy of single cells. *Photoacoustics*, 1(3), 49-53.
- Sugita, N., Hayashi, S., Hino, F. and Takanami, T. (2012). Palladium-catalyzed Kumada Coupling Reaction of bromoporphyrins with silylmethyl Grignard reagents: Preparation of silylmethyl-substituted porphyrins as a multipurpose synthon for fabrication of porphyrin systems. *The Journal of Organic Chemistry*, 77(23), 10488-10497.
- Sugita, N., Hayashi, S., Ishii, S. and Takanami, T. (2013). Palladium-Catalyzed Polyfluorophenylation of Porphyrins with Bis (polyfluorophenyl) zinc reagents. *Catalysts*, 3(4), 839-852.
- Suijkerbuijk, B. M. and Klein Gebbink, R. J. (2008). Merging porphyrins with organometallics: synthesis and applications. *Angewandte Chemie International Edition*, 47(39), 7396-7421.

- Sundholm, D. (2000). Density functional theory study of the electronic absorption spectrum of Mg-porphyrin and Mg-etiochlorophyll-I. *Chemical Physics Letters*, 317(3), 392-399.
- Suslick, K. S. and Watson, R. A. (1992). The photochemistry of chromium, manganese, and iron porphyrin complexes. *New Journal of Chemistry* 16, 633-642.
- Szabo, A. and Ostlund, N. S. (2012). *Modern quantum chemistry: introduction to advanced electronic structure theory*. New York: Courier Dover Publications.
- Takanami, T. (2013). Functionalization of Porphyrins through CC Bond Formation Reactions with Functional Group-Bearing Organometallic Reagents. *Heterocycles: an International Journal for Reviews and Communications in Heterocyclic Chemistry*, 87(8), 1659-1689.
- Takanami, T., Wakita, A., Sawaizumi, A., Iso, K., Onodera, H. and Suda, K. (2008). One-pot synthesis of meso-formylporphyrins by SNAr reaction of 5, 15-disubstituted porphyrins with (2-pyridyldimethylsilyl) methyllithium. *Organic Letters*, 10(4), 685-687.
- Taniguchi, M. and Lindsey, J. S. (2012). Enumeration of isomers of substituted tetrapyrrole macrocycles: from classical problems in biology to modern combinatorial libraries *Handbook of Porphyrin Science (Volume 23) With Applications to Chemistry, Physics, Materials Science, Engineering, Biology and Medicine* (pp. 1-80): World Scientific.
- Taniguchi, S., Hasegawa, H., Nishimura, M. and Takahashi, M. (1999). A Facile Route to Tripyrrane from 2, 5-Bis (hydroxymethyl) pyrrole and the Improved Synthesis of Porphine by the. *Synlett*, 1999(01), 73-74.
- Tapia, O. and Goscinski, O. (1975). Self-consistent reaction field theory of solvent effects. *Molecular Physics*, 29(6), 1653-1661.
- Tawada, Y., Tsuneda, T., Yanagisawa, S., Yanai, T. and Hirao, K. (2004). A long-range-corrected time-dependent density functional theory. *The Journal of Chemical Physics*, 120(18), 8425-8433.
- Taylor, G. A. and Tsutsui, M. (1975). Out-of-plane metalloporphyrin complexes. *Journal of Chemical Education*, 52(11), 715-720.
- Thies, S., Bornholdt, C., Köhler, F., Sönnichsen, F. D., Näther, C., Tuzek, F. and Herges, R. (2010). Coordination-Induced Spin Crossover (CISCO) through Axial Bonding of Substituted Pyridines to Nickel-Porphyrins:  $\sigma$ -Donor versus  $\pi$ -Acceptor Effects. *Chemistry-A European Journal*, 16(33), 10074-10083.
- Thudichum, J. (1867). Report Med. Off. Privy Council, 10, 152.

- Tirado-Rives, J. and Jorgensen, W. L. (2008). Performance of B3LYP density functional methods for a large set of organic molecules. *Journal of Chemical Theory and Computation*, 4(2), 297-306.
- To, W. P., Liu, Y., Lau, T. C. and Che, C. M. (2013). A Robust Palladium (II)–Porphyrin Complex as Catalyst for Visible Light Induced Oxidative C–H Functionalization. *Chemistry-A European Journal*, 19(18), 5654-5664.
- Tomasi, J., Mennucci, B. and Cammi, R. (2005). Quantum mechanical continuum solvation models. *Chemical Reviews*, 105(8), 2999-3094.
- Tovmasyan, A., Sheng, H., Weitner, T., Arulpragasam, A., Lu, M., Warner, D. S. . . . Batinic-Haberle, I. (2013). Design, mechanism of action, bioavailability and therapeutic effects of Mn porphyrin-based redox modulators. *Medical Principles and Practice*, 22(2), 103-130.
- Trova, M. P., Gauuan, P. J. F., Pechulis, A. D., Bubbs, S. M., Bocckino, S. B., Crapo, J. D. and Day, B. J. (2003). Superoxide dismutase mimetics. Part 2: synthesis and structure–Activity relationship of glyoxylate-and glyoxamide-Derived metalloporphyrins. *Bioorganic and Medicinal Chemistry*, 11(13), 2695-2707.
- Tsuchiya, S. and Seno, M. (1989). Novel synthetic method of phenol from benzene catalysed by perfluorinated hemin. *Chemistry Letters*, 18(2), 263-266.
- Tsutsui, M. and Hsung, C. (1973). A technetium porphyrin:  $\mu$ -[mesoporphyrin ix dimethyl esterato] bis [tricarbonyltechnetium (i)]. *Chemistry Letters*, 2(9), 941-942.
- Tsutsui, M., Ostfeld, D. and Hoffman, L. M. (1971). Unusual metalloporphyrins. VI. Metal shuttling between imidazole nitrogens in a ruthenium porphyrin imidazole. *Journal of the American Chemical Society*, 93(7), 1820-1823.
- Ullrich, C. A. (2011). *Time-dependent density-functional theory: concepts and applications*. Oxford: Oxford University Press.
- Urbani, M., Grätzel, M., Nazeeruddin, M. K. and Torres, T. (2014). Meso-substituted porphyrins for dye-sensitized solar cells. *Chemical Reviews*, 114(24), 12330-12396.
- Valeur, B. and Berberan-Santos, M. N. (2012). *Molecular fluorescence: principles and applications*. USA; John Wiley & Sons.
- Valicsek, Z., Eller, G. and Horváth, O. (2012). Equilibrium, photophysical and photochemical examination of anionic lanthanum (III) mono-and bisporphyrins: the effects of the out-of-plane structure. *Dalton Transactions*, 41(42), 13120-13131.

- Valicsek, Z. and Horváth, O. (2007). Formation, photophysics and photochemistry of thallium (III) 5, 10, 15, 20-tetrakis (4-sulphonatophenyl) porphyrin: New supports of typical sitting-atop features. *Journal of Photochemistry and Photobiology A: Chemistry*, 186(1), 1-7.
- Valicsek, Z. and Horváth, O. (2013). Application of the electronic spectra of porphyrins for analytical purposes: the effects of metal ions and structural distortions. *Microchemical Journal*, 107, 47-62.
- Valicsek, Z., Horváth, O. and Patonay, K. (2011). Formation, photophysical and photochemical properties of water-soluble bismuth (III) porphyrins: The role of the charge and structure. *Journal of Photochemistry and Photobiology A: Chemistry*, 226(1), 23-35.
- Valicsek, Z., Lendvay, G. r. and Horváth, O. (2008). Equilibrium, photophysical, photochemical, and quantum chemical examination of anionic mercury (II) mono-and bisporphyrins. *The Journal of Physical Chemistry B*, 112(46), 14509-14524.
- Van Gisbergen, S., Rosa, A., Ricciardi, G. and Baerends, E. (1999). Time-dependent density functional calculations on the electronic absorption spectrum of free base porph. *The Journal of Chemical Physics*, 111(6), 2499-2506.
- Vasil'ev, V. and Lobanovskaya, A. (2015). Production of hydrogen in a ternary photocatalytic system based on water-soluble palladium (II) porphyrin. *Russian Journal of Applied Chemistry*, 88(2), 283-288.
- Vaz, B., Alvarez, R., Nieto, M., Paniello, A. I. and de Lera, A. R. (2001). Suzuki cross-coupling of meso-dibromoporphyrins for the synthesis of functionalized A 2 B 2 porphyrins. *Tetrahedron Letters*, 42(42), 7409-7412.
- Vicente, M. (2001). Porphyrin-based sensitizers in the detection and treatment of cancer: recent progress. *Current Medicinal Chemistry-Anti-Cancer Agents*, 1(2), 175-194.
- Vicente, M. G. H. and Smith, K. M. (1990). Functionalization of alkyl substituents in octaalkylporphyrins. *Synlett*, 1990(10), 579-581.
- Vicente, M. G. H. and Smith, K. M. (1991). Vilsmeier reactions of porphyrins and chlorins with 3-(dimethylamino) acrolein to give meso-(2-formylvinyl) porphyrins: New syntheses of benzochlorins, benzoisobacteriochlorins, and benzobacteriochlorins and reductive coupling of porphyrins and chlorins using low-valent titanium complexes. *The Journal of Organic Chemistry*, 56(14), 4407-4418.

- Vydrov, O. A., Scuseria, G. E. and Perdew, J. P. (2007). Tests of functionals for systems with fractional electron number. *The Journal of Chemical Physics*, 126(15), 154109(1-9).
- Wang, B., Su, J. L., Amirian, J., Litovsky, S. H., Smalling, R. and Emelianov, S. (2010). Detection of lipid in atherosclerotic vessels using ultrasound-guided spectroscopic intravascular photoacoustic imaging. *Optics Express*, 18(5), 4889-4897.
- Wang, B., Zhao, Q., Barkey, N. M., Morse, D. L. and Jiang, H. (2012). Photoacoustic tomography and fluorescence molecular tomography: A comparative study based on indocyanine green. *Medical Physics*, 39(5), 2512-2517.
- Wang, C.-B. and Chang, C. (1979). A Convenient Synthesis of Pyrrole Precursors for Octaalkylporphyrins. *Synthesis*, 1979(07), 548-549.
- Wang, H., Zhou, D., Shen, S., Wan, J., Zheng, X., Yu, L. and Phillips, D. L. (2014). The photocatalytic activity and degradation mechanism of methylene blue over copper (ii) tetra (4-carboxyphenyl) porphyrin sensitized TiO<sub>2</sub> under visible light irradiation. *Rsc Advances*, 4(55), 28978-28986.
- Wang, L., Yang, P.-P., Zhao, X.-X. and Wang, H. (2016). Self-assembled nanomaterials for photoacoustic imaging. *Nanoscale*, 8(5), 2488-2509.
- Wang, L. V. and Hu, S. (2012). Photoacoustic tomography: in vivo imaging from organelles to organs. *Science*, 335(6075), 1458-1462.
- Wang, R., Brugh, A. M., Rawson, J., Therien, M. J. and Forbes, M. D. E. (2017). Alkyne-Bridged Multi[Copper(II) Porphyrin] Structures: Nuances of Orbital Symmetry in Long-Range, Through-Bond Mediated, Isotropic Spin Exchange Interactions. *Journal of the American Chemical Society*, 139(29), 9759-9762.
- Wang, T., Nandy, S., Salehi, H. S., Kumavor, P. D. and Zhu, Q. (2014). A low-cost photoacoustic microscopy system with a laser diode excitation. *Biomedical Optics Express*, 5(9), 3053-3058.
- Wang, T., Yang, Y., Alqasemi, U., Kumavor, P. D., Wang, X., Sanders, M. . . . Zhu, Q. (2013). Characterization of ovarian tissue based on quantitative analysis of photoacoustic microscopy images. *Biomedical Optics Express*, 4(12), 2763-2768.
- Weigend, F., Köhn, A. and Hättig, C. (2002). Efficient use of the correlation consistent basis sets in resolution of the identity MP2 calculations. *The Journal of Chemical Physics*, 116(8), 3175-3183.
- Weng, Y.-Q., Teng, Y.-L., Yue, F., Zhong, Y.-R. and Ye, B.-H. (2007). A new selective fluorescent chemosensor for Cu (II) ion based on zinc porphyrin-dipyridylamino. *Inorganic Chemistry Communications*, 10(4), 443-446.

- Weng, Y.-Q., Yue, F., Zhong, Y.-R. and Ye, B.-H. (2007). A Copper (II) Ion-Selective On– Off-Type Fluoroionophore Based on Zinc Porphyrin– Dipyritylamino. *Inorganic Chemistry*, 46(19), 7749-7755.
- Wiehe, A., Ryppa, C. and Senge, M. O. (2002). A practical synthesis of meso-monosubstituted,  $\beta$ -unsubstituted porphyrins. *Organic Letters*, 4(22), 3807-3809.
- Wijesekera, T., Dupre, D., Cader, M. S. and Dolphin, D. (1996). Porphyrins bearing halogens at the meso-phenyl and -pyrrolic positions: synthesis and spectral properties. *Bulletin-Societe Chimique De France-All Sections*, 133, 765-776.
- Wijesekera, T., Matsumoto, A., Dolphin, D. and Lexa, D. (1990). Perchlorinated and Highly Chlorinated meso-Tetraphenylporphyrins. *Angewandte Chemie International Edition*, 29(9), 1028-1030.
- Wijesekera, T. P. and Dolphin, D. (1990). 1-Bromo-19-methylbiladienes-ac; Useful precursors to porphyrins. *Synlett*, 1990(05), 235-244.
- Wijesekera, T. P. and Dolphin, D. (1994). Synthetic aspects of porphyrin and metalloporphyrin chemistry. *Metalloporphyrins in Catalytic Oxidations*, 193-239.
- Willard, H. H., Merritt Jr, L. L., Dean, J. A. and Settle Jr, F. A. (1988). *Instrumental methods of analysis*. New York: Wadsworth Publishing Company.
- Williams, A. T. R., Winfield, S. A. and Miller, J. N. (1983). Relative fluorescence quantum yields using a computer-controlled luminescence spectrometer. *Analyst*, 108(1290), 1067-1071.
- Willstätter, R. and Stoll, A. (1913). Die wirkungen der chlorophyllase. In: R. Willstätter and A. Stoll (eds.), *Untersuchungen ueber chlorophyll* (pp. 172-187). Berlin: Springer.
- Wolfram, K. and Holthausen, M. (2001). *A Chemist's Guide to Density Functional Theory*. Weinheim: Wiley–VCH Verlag, GmbH.
- Wróbel, D. and Graja, A. (2006). Modification of electronic structure in supramolecular fullerene–porphyrin systems studied by fluorescence, photoacoustic and photothermal spectroscopy. *Journal of Photochemistry and Photobiology A: Chemistry*, 183(1), 79-88.
- Wu, G.-Z., Gan, W.-X. and Leung, H.-K. (1991). Photophysical properties of meso-substituted octaethylporphines and their zinc complexes. *Journal of the Chemical Society, Faraday Transactions*, 87(18), 2933-2937.



- Wu, J. I., Fernández, I. and Schleyer, P. v. R. (2012). Description of aromaticity in porphyrinoids. *Journal of the American Chemical Society*, 135(1), 315-321.
- Xiang, L., Wang, B., Ji, L. and Jiang, H. (2013). 4-D photoacoustic tomography. *Scientific Reports*, 3, 1113.
- Xu, M. and Wang, L. V. (2006). Photoacoustic imaging in biomedicine. *Review of Scientific Instruments*, 77(4), 041101(1-22).
- Yamaguchi, T., Kimura, T., Matsuda, H. and Aida, T. (2004). Macroscopic Spinning Chirality Memorized in Spin-Coated Films of Spatially Designed Dendritic Zinc Porphyrin J-Aggregates. *Angewandte Chemie*, 116(46), 6510-6515.
- Yanai, T., Tew, D. P. and Handy, N. C. (2004). A new hybrid exchange–correlation functional using the Coulomb-attenuating method (CAM-B3LYP). *Chemical Physics Letters*, 393(1), 51-57.
- Yao, J., Wang, L., Yang, J.-M., Maslov, K. I., Wong, T. T., Li, L. . . . Wang, L. V. (2015). High-speed label-free functional photoacoustic microscopy of mouse brain in action. *Nature Methods*, 12(5), 407-410.
- Yasin, A., Jose, R. and Yusoff, M. M. (2015). Predicting larger absorption cross-section in porphyrin dyes using DFT calculations. *Journal of Porphyrins and Phthalocyanines*, 19(12), 1270-1278.
- Yeh, C.-Y., Miller, S. E., Carpenter, S. D. and Nocera, D. G. (2001). Structurally homologous  $\beta$ - and meso-amidinium porphyrins. *Inorganic Chemistry*, 40(14), 3643-3646.
- Yella, A., Lee, H.-W., Tsao, H. N., Yi, C., Chandiran, A. K., Nazeeruddin, M. K. . . . Grätzel, M. (2011). Porphyrin-sensitized solar cells with cobalt (II/III)–based redox electrolyte exceed 12 percent efficiency. *Science*, 334(6056), 629-634.
- Yelleswarapu, C. S. and Kothapalli, S.-R. (2010). Nonlinear photoacoustics for measuring the nonlinear optical absorption coefficient. *Optics Express*, 18(9), 9020-9025.
- Youngblood, W. J., Gryko, D. T., Lammi, R. K., Bocian, D. F., Holten, D. and Lindsey, J. S. (2002). Glaser-mediated synthesis and photophysical characterization of diphenylbutadiyne-linked porphyrin dyads. *The Journal of Organic Chemistry*, 67(7), 2111-2117.
- Yuan, Z., Wu, C., Zhao, H. and Jiang, H. (2005). Imaging of small nanoparticle-containing objects by finite-element-based photoacoustic tomography. *Optics Letters*, 30(22), 3054-3056.

- Zackrisson, S., van de Ven, S. and Gambhir, S. (2014). Light in and sound out: emerging translational strategies for photoacoustic imaging. *Cancer Research*, 74(4), 979-1004.
- Zems, Y., Moiseev, A. G. and Perepichka, D. F. (2013). Convenient synthesis of a highly soluble and stable phosphorescent platinum porphyrin dye. *Organic Letters*, 15(20), 5330-5333.
- Zhan, H., Lamare, S., Ng, A., Kenny, T., Guernon, H., Chan, W.-K. . . . Wong, W.-Y. (2011). Synthesis and photovoltaic properties of new metalloporphyrin-containing polyplatinyne polymers. *Macromolecules*, 44(13), 5155-5167.
- Zhang, D., Qi, G. B., Zhao, Y. X., Qiao, S. L., Yang, C. and Wang, H. (2015). In Situ Formation of Nanofibers from Purpurin18-Peptide Conjugates and the Assembly Induced Retention Effect in Tumor Sites. *Advanced Materials*, 27(40), 6125-6130.
- Zhang, X., Zhang, Y. and Jiang, J. (2005). Infrared spectra of metal-free, N', N-dideuterio, and magnesium porphyrins: density functional calculations. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 61(11), 2576-2583.
- Zhang, Y., Cai, X., Wang, Y., Zhang, C., Li, L., Choi, S. W. . . . Xia, Y. (2011). Noninvasive photoacoustic microscopy of living cells in two and three dimensions through enhancement by a metabolite dye. *Angewandte Chemie*, 123(32), 7497-7501.
- Zhao, Y. and Truhlar, D. G. (2006). Density functional for spectroscopy: no long-range self-interaction error, good performance for Rydberg and charge-transfer states, and better performance on average than B3LYP for ground states. *The Journal of Physical Chemistry A*, 110(49), 13126-13130.
- Zhao, Y. and Truhlar, D. G. (2008). The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals. *Theoretical Chemistry Accounts: Theory, Computation, and Modeling (Theoretica Chimica Acta)*, 120(1), 215-241.
- Zhou, Y., Liang, X. and Dai, Z. (2016). Porphyrin-loaded nanoparticles for cancer theranostics. *Nanoscale*, 8(25), 12394-12405.
- Zhu, W., Santic, M., Ou, Z., Santic, P. J., McDonald, J. A., Brotherhood, P. R. . . . Kadish, K. M. (2009). Electrochemistry and Spectroelectrochemistry of  $\beta$ ,  $\beta'$ -Fused Quinoxalinoporphyrins and Related Extended Bis-porphyrins with Co (III), Co (II), and Co (I) Central Metal Ions. *Inorganic Chemistry*, 49(3), 1027-1038.